

A strongly hyperbolic and regular reduction of Einstein's equations for axisymmetric spacetimes

Oliver Rinne* and John M Stewart†

*Department of Applied Mathematics and Theoretical Physics,
Centre for Mathematical Sciences, Wilberforce Road,
Cambridge CB3 0WA, United Kingdom*

(Dated: 9 February 2005)

This paper is concerned exclusively with axisymmetric spacetimes. We want to develop reductions of Einstein's equations which are suitable for numerical evolutions. We first make a Kaluza-Klein type dimensional reduction followed by an ADM reduction on the Lorentzian 3-space, the (2+1)+1 formalism. We include also the Z4 extension of Einstein's equations adapted to this formalism. Our gauge choice is based on a generalized harmonic gauge condition. We consider vacuum and perfect fluid sources.

We use these ingredients to construct a strongly hyperbolic first-order evolution system and exhibit its characteristic structure. This enables us to construct constraint-preserving stable outer boundary conditions. We use cylindrical polar coordinates and so we provide a careful discussion of the coordinate singularity on axis. By choosing our dependent variables appropriately we are able to produce an evolution system in which each and every term is manifestly regular on axis.

PACS numbers: 04.25.Dm, 04.20.Ex, 04.20.Cv

Keywords: Numerical relativity, axisymmetric spacetimes, regularity conditions, (2+1)+1 formalism, Z4 system, constraint-preserving boundary conditions, initial boundary value problem

I. INTRODUCTION

In the early days of numerical relativity attention focussed on spherically symmetric spacetimes. The next simplest subject would have been axisymmetric spacetimes, but very little progress was made in this area because, in polar coordinates adapted to the symmetry, there is a coordinate singularity on axis. Many attempts to deal with this proved unsuccessful, most evolutions became unstable and attention quickly turned, with the rapidly increasing capacity and speed of hardware, to spacetimes without symmetries. For a comprehensive review see e.g., [13].

However axisymmetric situations arise frequently in astrophysics and deserve further study. The first question is, obviously, why were the evolutions unstable? Suppose we assume that in a neighbourhood of the axis the spacetime is regular so that we can introduce locally cartesian coordinates (t, z, x, y) with the axis given by $x = y = 0$. Then the Killing vector is $\xi = -y\partial/\partial x + x\partial/\partial y$, and we say that M is *axisymmetric* if $\mathcal{L}_\xi M = 0$. We say that a tensor field M is *regular on axis* if its components with respect to this chart can be developed as a Taylor series in x and y in a neighbourhood of the axis¹. We may introduce polar coordinates $r = \sqrt{x^2 + y^2}$ and $\varphi = \arctan(y/x)$, so that the Killing vector is $\xi = (\partial/\partial\varphi)$. Because the transformation between cartesian coordinates and polar coordinates (t, z, r, φ) is singular on axis², axisymmetric tensor fields which are regular on axis may take strange forms when expressed in polar coordinates. For example if $M_{\alpha\beta}$ is symmetric and has these properties then

$$M_{\alpha\beta} = \begin{pmatrix} A & B & rD & r^2F \\ B & C & rE & r^2G \\ rD & rE & H + r^2J & r^3K \\ r^2F & r^2G & r^3K & r^2(H - r^2J) \end{pmatrix}, \quad (1)$$

where $A, B, \dots, K$ are functions of t, z and r^2 . (See appendix A for this and related results.) In particular both the metric and its first time-derivative must take this form. If the evolution scheme fails to preserve precisely the indicated r -dependencies then M becomes irregular on axis and instability is inevitable.

*Electronic address: o.rinne@damtp.cam.ac.uk

†Electronic address: j.m.stewart@damtp.cam.ac.uk

¹ For numerical purposes analytic and regular are synonymous.

² Note that the cartesian expression for the Killing vector ξ vanishes on axis, whereas the polar form does not.

Clearly the problems lie mainly in the last row and column of (1). In section II we outline a reduction of Einstein's equations in spacetime $\mathcal{M}$ to a 3-dimensional Lorentzian manifold $\mathcal{N}$ with chart (t, z, r) , effectively removing the last row/column. (The relation between M_{rr} and $M_{\varphi\varphi}$ still needs careful treatment.) Given a solution of the reduced equations in $\mathcal{N}$ one can recover the solution in $\mathcal{M}$. However there is no need to do this. All axisymmetric quantities of physical interest in $\mathcal{M}$ have their counterparts in $\mathcal{N}$, and so one might as well work in $\mathcal{N}$. Because that manifold is Lorentzian we can perform an ADM decomposition. The details of this $(2+1)+1$ scheme, with an arbitrary axisymmetric matter source, are given in section II.

In the $(2+1)+1$ system we have a spatial foliation with lapse function α , shift vector η^A , and spatial 2-metric H_{AB} . (Here $A, B, \dots$ range over r and z .) Since every 2-metric is conformally flat it is tempting to set $H_{rz} = 0$, $H_{rr} = H_{zz} = \psi^4$ say. The constraint equations imply elliptic equations for ψ and the shift vector components, and a quasi-maximal slicing condition supplies an elliptic equation for α . Although elliptic equations can become computationally expensive, multigrid techniques, if applicable, can reduce significantly the cost. The same choice was made by Choptuik et al [5]. Like those authors we encountered several difficulties, the most severe being that in near-critical collapse situations the discretized version of the energy constraint equation for ψ failed to remain diagonally dominant so that the multigrid algorithm failed to converge. Garfinkle et al [10] used a conjugate gradient method which gets round this particular problem. If instead we used an evolution equation for ψ there appeared to be a serious violation of the energy constraint.

We decided therefore to look at the other extreme, to minimize the number of elliptic equations solved. There are many, apparently different, ways of doing this, but we chose to implement the Z4 extension of Einstein's equation suggested by Bona et al [2], [3], but adapted to the $(2+1)+1$ scheme. The Z $(2+1)+1$ scheme is outlined in section III. There the energy and momentum constraints as well as the Geroch constraint (which arises in the $(2+1)+1$ decomposition) are converted into evolution equations for the new variables.

Pursuing this philosophy we would like to impose dynamical gauge conditions, which are discussed in section IV. These are generalizations of the harmonic gauge condition. However cylindrical polar coordinates are not harmonic, and this means that we can only do this for the lapse function. For simplicity we assume that the shift vector vanishes.

We now have a system of pure evolution equations and it is highly desirable to cast them into first order form. This is carried out in section V where we express them in conservation form with sources. We show in section VI that this system is *strongly hyperbolic* and we write down the conversions between conserved and characteristic variables. The characteristic speeds (divided by the lapse function α) are 0, ± 1 and $\pm\sqrt{f}$ where f is a constant parameter in the lapse evolution equation. (The choice $f = 1$ gives a harmonic time coordinate and a *symmetric hyperbolic* system.)

Knowing the characteristic structure enables us to formulate constraint-preserving boundary conditions for the initial-boundary value problem in section VII.

In section VIII we return to our starting point, regularity on axis. By a judicious choice of new dependent variables (which does not affect the hyperbolicity) we can write our first order strongly hyperbolic system in a form where each and every term is manifestly regular on axis.

Finally we discuss some of the implementation possibilities and applications of our scheme in section IX.

II. THE $(2+1)+1$ FORMALISM

Spacetime is assumed to be a four-dimensional manifold $(\mathcal{M}, g_{\alpha\beta})$ with signature $(-+++)$ and a preferred polar coordinate chart (t, r, z, φ) . Axisymmetry means that there is an everywhere spacelike Killing vector field $\xi = \partial/\partial\varphi$ with closed orbits. The idea is to perform first a Kaluza-Klein-like reduction from $\mathcal{M}$ to the Lorentzian three-dimensional manifold $\mathcal{N}$ formed by the trajectories of the Killing vector. (This was carried out for vacuum spacetimes by Geroch [11]. Nakamura et al [14, 15] extended the reduction to include general matter sources and then considered the case of a perfect fluid. Choptuik et al [5] added a massless scalar field source, albeit without rotation. The current analysis includes arbitrary sources and rotation.) After this an ADM-like reduction is applied to $\mathcal{N}$. We shall follow the notation of Nakamura et al. [14, 15], with some, clearly stated, changes. In the following, Greek indices range over t, r, z, φ , lower-case Latin indices over t, r, z , and upper-case Latin indices over r, z .

Tensor fields $M^{\alpha\cdots\beta\cdots}$ in $\mathcal{M}$ which are both orthogonal to ξ and are axisymmetric, i.e.,

$$M^{\alpha\cdots\beta\cdots}\xi_\beta = M^{\alpha\cdots\beta\cdots}\xi_\alpha = \dots = 0, \quad \mathcal{L}_\xi M^{\alpha\cdots\beta\cdots} = 0$$

are said to be tensor fields *in* $\mathcal{N}$. Some basic tensor fields in $\mathcal{N}$ are the norm of the Killing vector

$$\lambda^2 = g_{\alpha\beta}\xi^\alpha\xi^\beta > 0,$$

the (Lorentzian) metric in $\mathcal{N}$,

$$h_{\alpha\beta} = g_{\alpha\beta} - \lambda^{-2}\xi_\alpha\xi_\beta,$$

the Levi-Civita tensor

$$\epsilon_{\alpha\beta\gamma} = \lambda^{-1} \epsilon_{\alpha\beta\gamma\delta} \xi^\delta,$$

and the “twist vector”

$$\omega_\alpha = \epsilon_{\alpha\beta\gamma\delta} \xi^\beta \nabla^\gamma \xi^\delta, \quad (2)$$

where ∇ is the covariant derivative of $g_{\alpha\beta}$. Note that neither $\xi^\alpha = \delta^\alpha_\varphi$ nor $\xi_\alpha = g_{\alpha\varphi}$ is in $\mathcal{N}$. However if a solution of the equations in $\mathcal{N}$ is given then the solution in $\mathcal{M}$ can be reconstructed [11]. We shall argue that a reconstruction is not necessary. All physically relevant quantities in $\mathcal{M}$ have their counterparts in $\mathcal{N}$.

We define the projections of the energy-momentum tensor as follows,

$$\begin{aligned} \tau &= \lambda^{-2} \xi^\gamma \xi^\delta T_{\gamma\delta}, \\ \tau_\alpha &= \lambda^{-2} h_\alpha{}^\gamma \xi^\delta T_{\gamma\delta}, \\ \tau_{\alpha\beta} &= h_\alpha{}^\gamma h_\beta{}^\delta T_{\gamma\delta}. \end{aligned} \quad (3)$$

Note that near the axis $\lambda = O(r)$, and the factors of λ are included in (3) to ensure that the projections are either $O(1)$ or $O(r)$ near the axis, so that regularity conditions (cf. section VIII) are easily imposed.

Following the standard ADM procedure, the three-dimensional manifold $\mathcal{N}$ is foliated into two-dimensional spacelike hypersurfaces $\Sigma(t)$ with unit normal $n_a = -\alpha \partial_a t$ and intrinsic metric

$$H_{ab} = h_{ab} + n_a n_b.$$

The line element in $\mathcal{N}$ takes the form

$$ds^2 = -\alpha^2 dt^2 + H_{AB} (dx^A + \eta^A dt) (dx^B + \eta^B dt),$$

where α is the lapse function and η^A is the shift vector. The extrinsic curvature is given by

$$\chi_{ab} = -H_a{}^c H_b{}^d n_{(c|d)} = -\frac{1}{2} \mathcal{L}_n H_{ab},$$

where $|$ denotes the covariant derivative compatible with h_{ab} . The trace of the extrinsic curvature is abbreviated as $\chi = \chi_A{}^A$. Further variables defined in each $\Sigma(t)$ are the alternating symbol

$$\epsilon_{AB} = n^c \epsilon_{cAB},$$

the φ -component of the extrinsic curvature,

$$K_\varphi{}^\varphi = -\lambda^{-1} n^a \lambda_{,a},$$

and the rotational degrees of freedom,

$$\begin{aligned} E^A &= \lambda^{-3} \epsilon^{Ab} \omega_b, \\ B^\varphi &= \lambda^{-3} n_a \omega^a. \end{aligned}$$

Again, the last two definitions differ from those in [14, 15] by factors of λ , and we write B^φ with an upstairs index. The various projections of the energy-momentum tensor are

$$\begin{aligned} J^\varphi &= -n_a \tau^a, \\ S^A &= H_a{}^A \tau^a, \\ \rho_H &= n^a n^b \tau_{ab}, \\ J_A &= -H_A{}^a n^b \tau_{ab}, \end{aligned} \quad (4)$$

$$S_{AB} = H_A{}^a H_B{}^b \tau_{ab}, \quad (5)$$

and, of course τ defined in (3).

For convenience we also introduce

$$A_A \equiv \alpha^{-1} \alpha_{,A}, \quad L_A \equiv \lambda^{-1} \lambda_{,A}. \quad (6)$$

We can now write down the projections of Einstein's equations

$$G_{\alpha\beta} = \kappa T_{\alpha\beta},$$

which split into sets of elliptic constraint equations and hyperbolic evolution equations.

The constraint equations are $\mathcal{C} = \mathcal{C}_A = \mathcal{C}_\varphi = 0$, where the energy constraint is

$$\mathcal{C} \equiv \frac{1}{2}(\chi^2 - \chi_{AB}\chi^{AB} + {}^{(2)}R) - L^A{}_{||A} - L_A L^A + \chi K_\varphi{}^\varphi - \frac{1}{4}\lambda^2 (E_A E^A + B^{\varphi 2}) - \kappa \rho_H, \quad (7)$$

with $||$ denoting the covariant derivative of H_{AB} , ${}^{(2)}R_{AB}$ the Ricci tensor of H_{AB} and ${}^{(2)}R = {}^{(2)}R_A{}^A$ the Ricci scalar. The linear momentum constraints are

$$\mathcal{C}_A \equiv \chi_A{}^B{}_{||B} - (\chi + K_\varphi{}^\varphi)_{,A} + L^B \chi_{AB} - L_A K_\varphi{}^\varphi - \frac{1}{2}\lambda^2 B^\varphi \epsilon_{AB} E^B - \kappa J_A, \quad (8)$$

and the angular momentum or Geroch constraint is

$$\mathcal{C}_\varphi \equiv \frac{1}{2}E^A{}_{||A} + \frac{3}{2}L_A E^A - \kappa J^\varphi. \quad (9)$$

We shall express all the evolution equations in terms of the Lie derivative along the normal direction,

$$\mathcal{L}_n = \alpha^{-1}(\partial_t - \mathcal{L}_\eta).$$

The time evolution of the metric variables is given by

$$\mathcal{L}_n H_{AB} = -2\chi_{AB}, \quad (10)$$

$$\mathcal{L}_n \lambda = -\lambda K_\varphi{}^\varphi. \quad (11)$$

The evolution equations for the extrinsic curvature variables are

$$\begin{aligned} \mathcal{L}_n \chi_{AB} &= {}^{(2)}R_{AB} + (\chi + K_\varphi{}^\varphi)\chi_{AB} - 2\chi_A{}^C \chi_{CB} - A_{A||B} - A_A A_B - L_{A||B} - L_A L_B \\ &\quad - \frac{1}{2}\lambda^2 [\epsilon_{AC}\epsilon_{BD}E^C E^D - H_{AB}(E_C E^C - B^{\varphi 2})] \\ &\quad - \kappa [S_{AB} + \frac{1}{2}H_{AB}(\rho_H - S_C{}^C - \tau)], \end{aligned} \quad (12)$$

$$\begin{aligned} \mathcal{L}_n K_\varphi{}^\varphi &= K_\varphi{}^\varphi(\chi + K_\varphi{}^\varphi) - L^A{}_{||A} - L_A(L^A + A^A) - \frac{1}{2}\lambda^2 (E_A E^A - B^{\varphi 2}) \\ &\quad - \frac{1}{2}\kappa(\rho_H - S_A{}^A + \tau). \end{aligned} \quad (13)$$

The rotation variables evolve according to

$$\mathcal{L}_n E^A = (\chi + 3K_\varphi{}^\varphi)E^A + \epsilon^{AB}B^\varphi{}_{,B} + \epsilon^{AB}(A_B + 3L_B)B^\varphi - 2\kappa S^A, \quad (14)$$

$$\mathcal{L}_n B^\varphi = \chi B^\varphi + \epsilon^{AB}E_A A_B + \epsilon^{AB}E_{A||B}. \quad (15)$$

(The principal parts of these equations resemble the axisymmetric Maxwell equations, which justifies the notation. The Geroch constraint (9) is also a ‘‘Maxwell equation’’.) Energy-momentum conservation $\nabla_\alpha T^{\alpha\beta} = 0$ implies the following evolution equations for the matter variables: the energy conservation equation

$$\mathcal{L}_n \rho_H = -J^A{}_{||A} - J^A(2A_A + L_A) + \chi \rho_H + \chi_{AB}S^{AB} + K_\varphi{}^\varphi \tau + \lambda^2 E^A S_A, \quad (16)$$

the Euler equations

$$\begin{aligned} \mathcal{L}_n J_A &= -S_{AB}{}^{||B} + J_A(\chi + K_\varphi{}^\varphi) - S_{AB}(A^B + L^B) - A_A \rho_H + L_A \tau \\ &\quad + \lambda^2 (E_A J^\varphi + \epsilon_{AB}S^B B^\varphi), \end{aligned} \quad (17)$$

and angular momentum conservation

$$\mathcal{L}_n J^\varphi = -S^A{}_{||A} - S^A(A_A + 3L_A) + J^\varphi(\chi + 3K_\varphi{}^\varphi). \quad (18)$$

In many situations there may be a conserved particle number density N^α satisfying $\mathcal{L}_\xi N^\alpha = 0$. If we introduce the projections

$$\begin{aligned} \nu^\alpha &= h^\alpha{}_\beta N^\beta, \\ \Sigma^A &= H^A{}_b \nu^b, \\ \sigma &= -\nu^a n_a, \end{aligned}$$

then particle number conservation $\nabla_\alpha N^\alpha = 0$ implies the following evolution equation for σ ,

$$\mathcal{L}_n \sigma = -\Sigma^A{}_{||A} - \Sigma^A(A_A + L_A) + \sigma(\chi + K_\varphi{}^\varphi). \quad (19)$$

III. THE Z4 EXTENSION

Bona et al. [2] suggested the following covariant extension of the Einstein field equations:

$$R_{\alpha\beta} + 2\nabla_{(\alpha}Z_{\beta)} = \kappa (T_{\alpha\beta} - \frac{1}{2}Tg_{\alpha\beta}) , \quad (20)$$

which reduces to the Einstein equations if and only if $Z_\alpha = 0$.

We shall assume that Z_α shares the axisymmetry, i.e., $\mathcal{L}_\xi Z_\alpha = 0$. To apply the (2+1)+1 formalism to equations (20), it is convenient to rewrite them as Einstein's equations $G_{\alpha\beta} = \kappa \tilde{T}_{\alpha\beta}$ with a modified energy-momentum tensor

$$\tilde{T}_{\alpha\beta} = T_{\alpha\beta} - \frac{2}{\kappa} (\nabla_{(\alpha}Z_{\beta)} - \frac{1}{2}g_{\alpha\beta}\nabla_\gamma Z^\gamma) . \quad (21)$$

The Z covector is decomposed into parts parallel and orthogonal to the Killing vector ξ^α ,

$$Z^\varphi = \lambda^{-2}\xi^\alpha Z_\alpha , \quad Z_a = h_a{}^\alpha Z_\alpha ,$$

and further parallel and orthogonal to the timelike normal n_a ,

$$\theta = -n^a Z_a , \quad Z_A = H_A{}^a Z_a .$$

We now compute the modified matter variables corresponding to $\tilde{T}_{\alpha\beta}$, finding

$$\begin{aligned} \tilde{\tau} &= \tau + \kappa^{-1} [\mathcal{L}_n \theta + Z^A{}_{||A} + (A_A - L_A)Z^A + (K_\varphi{}^\varphi - \chi)\theta] , \\ \tilde{S}_A &= S_A + \kappa^{-1} [-Z^\varphi{}_{,A} + B^\varphi \epsilon_{AB} Z^B + E_A \theta] , \\ \tilde{J}^\varphi &= J^\varphi + \kappa^{-1} [\mathcal{L}_n Z^\varphi + E^A Z_A] , \\ \tilde{S}_{AB} &= S_{AB} + \kappa^{-1} [-2Z_{(A||B)} + 2\chi_{AB}\theta \\ &\quad + H_{AB} \{ \mathcal{L}_n \theta + Z^C{}_{||C} + (A_C + L_C)Z^C - (\chi + K_\varphi{}^\varphi)\theta \}] , \\ \tilde{J}_A &= J_A + \kappa^{-1} [\mathcal{L}_n Z_A - \theta_{,A} + 2\chi_{AB} Z^B + A_A \theta] , \\ \tilde{\rho}_H &= \rho_H + \kappa^{-1} [\mathcal{L}_n \theta - Z^A{}_{||A} + (A_A - L_A)Z^A + (\chi + K_\varphi{}^\varphi)\theta] . \end{aligned} \quad (22)$$

Inserting the above into the standard (2+1)+1 equations, we obtain the Z(2+1)+1 equations. Because of the Lie derivatives in (22), the constraints (7-9) become evolution equations for the Z covector,

$$\mathcal{L}_n \theta = C + (L_A - A_A)Z^A + Z^A{}_{||A} - (\chi + K_\varphi{}^\varphi)\theta , \quad (23)$$

$$\mathcal{L}_n Z_A = C_A - 2\chi_{AB} Z^B - A_A \theta + \theta_{,A} , \quad (24)$$

$$\mathcal{L}_n Z^\varphi = C_\varphi - E^A Z_A . \quad (25)$$

The main evolution equations are modified as follows,

$$\begin{aligned} \mathcal{L}_n \chi_{AB} &= \dots + 2Z_{(A||B)} - 2\chi_{AB}\theta , \\ \mathcal{L}_n K_\varphi{}^\varphi &= \dots + 2L_A Z^A - 2K_\varphi{}^\varphi \theta , \\ \mathcal{L}_n E^A &= \dots + 2Z^\varphi{}_{,A} - 2E^A \theta - 2B^\varphi \epsilon^{AB} Z_B , \end{aligned}$$

where $\dots$ denote the right-hand-sides of (12), (13) and (14), respectively. The remaining evolution equations are unchanged.

IV. DYNAMICAL GAUGE CONDITIONS

To complete our evolution formalism, we need to prescribe the gauge variables α and η^A . It is well-known [6, 8] that the Einstein equations can be written in symmetric hyperbolic form

$$g^{\gamma\delta} g_{\alpha\beta,\gamma\delta} \simeq 0$$

($\simeq$ denoting equality to principal parts) by adopting harmonic gauge, which for the Z4 extension of the field equations becomes

$$g^{\gamma\delta} x^\alpha_{;\gamma\delta} = 2Z^\alpha,$$

where we are treating x^α as a scalar field. It would therefore be desirable to choose this or a related gauge for the system at hand. However, these gauge conditions are incompatible with cylindrical polar coordinates; the wave operator contains a term $r^{-1}\partial_r$, and the right-hand-side of the evolution equation for the shift vector becomes singular on the axis. However, we can keep the time component of the harmonic gauge equation, which in (2+1)+1 language reads

$$\partial_t \alpha = -\alpha^2 (\chi + K_\varphi^\varphi - 2\theta).$$

This condition has been generalized to [3]

$$\partial_t \alpha = -f\alpha^2 (\chi + K_\varphi^\varphi - m\theta), \quad (26)$$

where f and m are free constant parameters.

We choose the shift vector to vanish: $\eta^A = 0$. Apart from simplifying the evolution equations, this choice also makes it much easier to set up outer boundary conditions because a variable shift would mean that the characteristic speeds could change sign at the boundary (see section VI). Our choice for the shift vector means that henceforth

$$\mathcal{L}_n = \alpha^{-1} \partial_t.$$

V. FIRST-ORDER REDUCTION

The Z(2+1)+1 equations are first-order in time but second-order in space. In order to analyze their hyperbolicity and to be able to treat them with standard numerical methods for hyperbolic conservation laws, we perform a first-order reduction. We focus on the geometry evolution equations here; for given geometry, the matter evolution equations form a separate strongly hyperbolic system (see appendix D).

To eliminate the second spatial derivatives, we introduce new variables for the first spatial derivatives of the two-metric,

$$D_{ABC} \equiv \frac{1}{2} \partial_A H_{BC},$$

and we regard A_A and L_A defined by (6) as independent variables. Evolution equations for these new first-order variables can be obtained from (10), (11) and (26) by commuting space and time derivatives:

$$\mathcal{L}_n D_{ABC} = -\chi_{BC,A}, \quad (27)$$

$$\mathcal{L}_n L_A = -K_{\varphi^\varphi,A}, \quad (28)$$

$$\mathcal{L}_n A_A = -f(\chi_{,A} + K_{\varphi^\varphi,A} - m\theta_{,A}). \quad (29)$$

The two independent traces of D_{ABC} are denoted by

$$D^I{}_A \equiv D_{AB}{}^B, \quad D^{II}{}_A \equiv D^B{}_{BA},$$

where indices are (formally, for D is not a tensor) raised and lowered with the 2-metric H_{AB} . The De Donder-Fock decomposition [6, 8] is used for the Ricci tensor,

$$\begin{aligned} {}^{(2)}R_{AB} = & -D^C{}_{AB,C} + 2D^{II}{}_{(A,B)} - D^I{}_{(A,B)} - 2D_{CAB}D^{II}{}^C \\ & - \Gamma_{CAB}(2D^{II}{}^C - D^I{}^C) + 4D_{CDA}D^{CD}{}_B - \Gamma_{ACD}\Gamma_B{}^{CD}, \end{aligned}$$

where of course the Christoffel symbols are given by

$$\Gamma_{ABC} = D_{CAB} + D_{BCA} - D_{ABC}. \quad (30)$$

It is then straightforward to write the Z(2+1)+1 equations in *conservation form*

$$\partial_t \mathbf{u} + [\alpha \mathcal{F}^D(\mathbf{u})]_{,D} = \alpha \mathcal{S}(\mathbf{u}). \quad (31)$$

Here, the set of conserved variables is

$$\mathbf{u} = (H_{AB}, \lambda, \alpha, D_{ABC}, L_A, A_A, \chi_{AB}, K_\varphi^\varphi, E^A, B^\varphi, \theta, Z_A, Z^\varphi)^T,$$

and the components of the flux vector are given by

$$\begin{aligned} \mathcal{F}^D_{H_{AB}} &= 0, \\ \mathcal{F}^D_\lambda &= 0, \\ \mathcal{F}^D_\alpha &= 0, \\ \mathcal{F}^D_{D_{ABC}} &= \delta_A^D \chi_{BC}, \\ \mathcal{F}^D_{L_A} &= \delta_A^D K_\varphi^\varphi, \\ \mathcal{F}^D_{A_A} &= \delta_A^D f(\chi + K_\varphi^\varphi - m\theta), \\ \mathcal{F}^D_{\chi_{AB}} &= D^D_{AB} - \delta_{(A}^D (2D^{II}_{B)} + 2Z_B) - D^I_{B)} - L_B - A_B), \\ \mathcal{F}^D_{K_\varphi^\varphi} &= L^D, \\ \mathcal{F}^D_{E^A} &= -2H^{AD} Z^\varphi - \epsilon^{AD} B^\varphi, \\ \mathcal{F}^D_{B^\varphi} &= -\epsilon^{AD} E_A, \\ \mathcal{F}^D_\theta &= D^{ID} - D^{IID} + L^D - Z^D, \\ \mathcal{F}^D_{Z_A} &= -\chi_A^D + \delta_A^D (\chi + K_\varphi^\varphi - \theta), \\ \mathcal{F}^D_{Z^\varphi} &= -\frac{1}{2} E^D. \end{aligned}$$

$\mathcal{S}(\mathbf{u})$ is a source term containing no derivatives,

$$\begin{aligned} \mathcal{S}_{H_{AB}} &= -2\chi_{AB}, \\ \mathcal{S}_\lambda &= -\lambda K_\varphi^\varphi, \\ \mathcal{S}_\alpha &= -f\alpha(\chi + K_\varphi^\varphi - m\theta), \\ \mathcal{S}_{D_{ABC}} &= 0, \\ \mathcal{S}_{L_A} &= 0, \\ \mathcal{S}_{A_A} &= 0, \\ \mathcal{S}_{\chi_{AB}} &= A_C D^C_{AB} + A_{(A} (-2D^{II}_{B)} + D^I_{B)} + L_B) + A_B - 2Z_B) \\ &\quad - L_A L_B - A_A A_B + \Gamma^C_{AB} (L_C + A_C) + \chi_{AB} (\chi + K_\varphi^\varphi) - 2\chi_A^C \chi_{CB} \\ &\quad - \frac{1}{2} \lambda^2 [\epsilon_{AC} \epsilon_{BD} E^C E^D - H_{AB} (E_C E^C - B^{\varphi 2})] \\ &\quad - \kappa [S_{AB} + \frac{1}{2} H_{AB} (\rho_H - S_C^C - \tau)] - \Gamma_{CAB} (2Z^C + 2D^{II\,C} - D^{I\,C}) \\ &\quad - 2\theta \chi_{AB} - 2D_{CAB} D^{II\,C} + 4D_{CDA} D^{CD}_B - \Gamma_{ACD} \Gamma_B^{CD}, \\ \mathcal{S}_{K_\varphi^\varphi} &= K_\varphi^\varphi (\chi + K_\varphi^\varphi) + L_A (2Z^A - L^A - D^{IA}) - \frac{1}{2} \lambda^2 (E_A E^A + B^{\varphi 2}) - 2K_\varphi^\varphi \theta \\ &\quad - \frac{1}{2} \kappa (\rho_H - S_C^C + \tau), \\ \mathcal{S}_{E^A} &= (\chi + 3K_\varphi^\varphi - 2\theta) E^A + \epsilon^{AB} B^\varphi (3L_B - 2Z_B + D^I_B) - 2\kappa S^A + (4D^{II\,A} - 2A^A) Z^\varphi, \\ \mathcal{S}_{B^\varphi} &= \chi B^\varphi + \epsilon^{AB} E_A D^I_B, \\ \mathcal{S}_\theta &= A_A (D^{IA} - D^{II\,A} + L^A - Z^A) + D_{ABC} D^{ABC} - \frac{1}{2} \Gamma_{ABC} \Gamma^{ABC} - \frac{1}{2} D^I_A D^{IA} \\ &\quad - L_A (L^A + D^{IA}) + \frac{1}{2} (\chi^2 - \chi_{AB} \chi^{AB}) + \chi K_\varphi^\varphi - \frac{1}{4} \lambda^2 (E_A E^A + B^{\varphi 2}) - \kappa \rho_H \\ &\quad + (L_A - A_A + D^I_A) Z^A - (\chi + K_\varphi^\varphi) \theta, \\ \mathcal{S}_{Z_A} &= -A_B \chi_A^B + A_A (\chi + K_\varphi^\varphi - \theta) + \chi_{AB} (D^{IB} + L^B - 2Z^B) - \Gamma^C_{AB} \chi_C^B \\ &\quad - L_A K_\varphi^\varphi - \frac{1}{2} \lambda^2 B^\varphi \epsilon_{AB} E^B - \kappa J_A - A_A \theta, \\ \mathcal{S}_{Z^\varphi} &= \frac{1}{2} E^A (D^I_A + 3L_A - 2Z_A - A_A) - \kappa J^\varphi. \end{aligned}$$

Note that H_{AB} , λ and α have vanishing fluxes and thus trivially propagate along the normal direction. The rotation variables $(E^A, B^\varphi, Z^\varphi)$ form a decoupled subsystem analogous to Maxwell's equations.

VI. HYPERBOLICITY

To investigate the hyperbolicity of the $Z(2+1)+1$ system, we pick a unit covector μ_A and define an orthogonal covector

$$\pi_A = \epsilon_{AB} \mu^B, \quad (32)$$

so that

$$\mu_A \mu^A = \pi_A \pi^A = 1, \quad \mu_A \pi^A = 0.$$

Thus (μ^A, π^A) form an orthonormal basis for $T(\Sigma)$. Projection along μ and π is denoted as

$$V^\parallel \equiv V^A \mu_A, \quad V^\perp \equiv V^A \pi_A. \quad (33)$$

It can be verified that the Jacobian matrix

$$B^\parallel \equiv \frac{\partial \mathcal{F}^\parallel}{\partial \mathbf{u}}$$

corresponding to the flux in the μ -direction is real, diagonalizable and has complete sets of left and right eigenvectors for arbitrary μ_A , i.e., the system is *strongly hyperbolic*. The characteristic speeds are³

$$\begin{aligned} \lambda_0 &= 0, \\ \lambda_1^\pm &= \pm 1 \quad (\text{speed of light}), \\ \lambda_f^\pm &= \pm \sqrt{f} \quad (\text{gauge speed}). \end{aligned}$$

For $f \leq 1$, all characteristic speeds are causal, for $f = 1$ (harmonic slicing) they are all physical.

The characteristic variables (left eigenvectors dotted into the conserved variables) are given by

- Normal modes (eigenvalue 0)

$$\begin{aligned} l_{0,1} &= fm(D_{\parallel\perp\perp} + L_\parallel - D_{\perp\perp\parallel} - Z_\parallel) - f(D_{\parallel\parallel\parallel} + D_{\parallel\perp\perp} + L_\parallel) + A_\parallel, \\ l_{0,2} &= fm(D_{\perp\parallel\parallel} + L_\perp - D_{\parallel\parallel\perp} - Z_\perp) - f(D_{\perp\parallel\parallel} + D_{\perp\perp\perp} + L_\perp) + A_\perp, \\ l_{0,3} &= D_{\perp\parallel\parallel}, \\ l_{0,4} &= D_{\perp\perp\parallel}, \\ l_{0,5} &= D_{\perp\perp\perp}, \\ l_{0,6} &= L_\perp, \\ l_{0,7} &= A_\perp, \end{aligned}$$

along with the zeroth-order variables H_{AB} , λ and α .

- Light cone modes (eigenvalue ± 1)

$$\begin{aligned} l_{1,1}^\pm &= \theta \pm (D_{\parallel\perp\perp} + L_\parallel - D_{\perp\perp\parallel} - Z_\parallel), \\ l_{1,2}^\pm &= \chi_{\parallel\perp} \pm \frac{1}{2}(A_\perp + D_{\perp\parallel\parallel} - D_{\perp\perp\perp} + L_\perp - 2Z_\perp), \\ l_{1,3}^\pm &= \chi_{\perp\perp} \pm D_{\parallel\perp\perp}, \\ l_{1,4}^\pm &= K_\varphi^\varphi \pm L_\parallel, \\ l_{1,5}^\pm &= E^\parallel \mp 2Z^\varphi, \\ l_{1,6}^\pm &= E^\perp \mp B^\varphi, \end{aligned}$$

³ Note that a factor of α has been taken out of the fluxes (31). Note also that for a nonzero shift vector η^A , a term $-\alpha^{-1}\eta^\parallel$ would have to be added to the above characteristic speeds.

- Lapse cone modes (eigenvalue $\pm\sqrt{f}$)

$$l_f^\pm = A_\parallel - \frac{f(m-2)}{f-1}(D_{\parallel\perp\perp} + L_\parallel - D_{\perp\perp\parallel} - Z_\parallel) \pm \sqrt{f} \left[\chi_{\parallel\parallel} + \chi_{\perp\perp} + K_\varphi^\varphi - \left(\frac{f(m-2)}{f-1} + 2 \right) \theta \right]. \quad (34)$$

The inverse transformation from characteristic to conserved variables is given in Appendix B.

In the special case $f = 1$, we must set $m = 2$ to enforce strong hyperbolicity, and the singular term $(m-2)/(f-1)$ in (34) is to be replaced with an arbitrary fixed constant (e.g. 0 for simplicity). In this case, the system is even *symmetric hyperbolic*, i.e. $B^\parallel$ is symmetrizable with a symmetrizer that is independent of the direction μ_A . This is not surprising because the choice of parameters $f = 1$, $m = 2$ in (26) corresponds to harmonic slicing. An energy for the system is given by

$$\begin{aligned} \mathcal{E} = & \chi_{AB}\chi^{AB} + \lambda_{ABC}\lambda^{ABC} + (K_\varphi^\varphi + \chi - 2\theta)^2 + A_A A^A \\ & + V_A V^A + K_\varphi^{\varphi^2} + L_A L^A + E_A E^A + B^{\varphi^2} + 4Z^{\varphi^2}, \end{aligned}$$

where

$$\begin{aligned} V_A &\equiv A_A + D^I{}_A + L_A - 2D^{II}{}_A - 2Z_A, \\ \lambda^A{}_{BC} &\equiv D^A{}_{BC} + \delta^A_{(B} V_{C)}. \end{aligned}$$

$\mathcal{E}$ is positive definite and its time derivative is (considering principal parts only) a total divergence.

VII. CONSTRAINT-PRESERVING BOUNDARY CONDITIONS

In the $Z(2+1)+1$ formalism, the standard $(2+1)+1$ constraints (7–9) are replaced with the *algebraic* constraints

$$\theta = Z_A = Z^\varphi = 0. \quad (35)$$

If those hold at all times, the $(2+1)+1$ constraints are automatically satisfied by virtue of the evolution equations for the Z vector (23–25). On the initial time level, one imposes (35) and solves the $(2+1)+1$ constraints (7–9) so that both the Z vector and its time derivative vanish. In addition, the outer boundary conditions have to be chosen so that no constraint-violating modes propagate into the computational domain at any time.

A propagation system for (35) follows from the contracted Bianchi identities, which imply $\nabla_\alpha \tilde{T}^{\alpha\beta} = 0$. If we impose the standard matter evolution equations $\nabla_\alpha T^{\alpha\beta} = 0$ in addition, (21) yields the following homogeneous wave equation for the Z covector:

$$\nabla_\beta \nabla^\beta Z_\alpha + R_{\alpha\beta} Z^\beta = 0. \quad (36)$$

To obtain the $(2+1)+1$ reduction of (36), we take an additional time derivative of the evolution equations for the Z vector (23–25). To principal parts, we have

$$\begin{aligned} \mathcal{L}_n^2 \theta - \partial_B \partial^B \theta &\simeq 0, \\ \mathcal{L}_n^2 Z_A - \partial_B \partial^B Z_A &\simeq \frac{1}{2} \partial_B [\partial_A (A^B + D^I{}^B + L^B) - \partial^B (A_A + D_{IA} + L_A) \\ &\quad - 2(\partial_C D^{BC}{}_A - \partial^B D^{II}{}_A)] , \\ \mathcal{L}_n^2 Z^\varphi - \partial_B \partial^B Z^\varphi &\simeq 0. \end{aligned} \quad (37)$$

Note that the right-hand-sides are zero if the *ordering constraints*

$$\begin{aligned} \partial_D D_{ABC} &= \partial_A D_{DBC}, \\ \partial_B L_A &= \partial_A L_B, \\ \partial_B A_A &= \partial_A A_B \end{aligned} \quad (38)$$

are satisfied, which corresponds to commuting of partial derivatives in the definition of the first-order variables. Although this holds analytically, it may not obtain numerically, and so the terms have to be included in the subsidiary system (37).

We choose an orthonormal basis μ^A, π^A such that μ^A is normal to the outer boundary under consideration. Indices contracted with μ^A (π^A) are denoted by $\parallel$ ($\perp$). Equations (37) decompose into

$$\begin{aligned}
\mathcal{L}_n^2 \theta - \partial_{\parallel}^2 \theta - \partial_{\perp}^2 \theta &\simeq 0, \\
\mathcal{L}_n^2 Z_{\parallel} - \partial_{\parallel}^2 Z_{\parallel} - \partial_{\perp}^2 Z_{\parallel} &\simeq \partial_{\parallel}^2 D_{\perp\perp\parallel} \\
&\quad + \frac{1}{2} \partial_{\parallel} \partial_{\perp} (A_{\perp} - D_{\perp\parallel\parallel} + D_{\perp\perp\perp} - 2D_{\parallel\parallel\perp} + L_{\perp}) \\
&\quad + \frac{1}{2} \partial_{\perp}^2 (-A_{\parallel} + D_{\parallel\parallel\parallel} - D_{\parallel\perp\perp} - L_{\parallel}), \\
\mathcal{L}_n^2 Z_{\perp} - \partial_{\parallel}^2 Z_{\perp} - \partial_{\perp}^2 Z_{\perp} &\simeq \frac{1}{2} \partial_{\parallel}^2 (-A_{\perp} - D_{\perp\parallel\parallel} + D_{\perp\perp\perp} - L_{\perp}) \\
&\quad + \frac{1}{2} \partial_{\parallel} \partial_{\perp} (A_{\parallel} + D_{\parallel\parallel\parallel} - D_{\parallel\perp\perp} - 2D_{\perp\perp\parallel} + L_{\parallel}) \\
&\quad + \partial_{\perp}^2 D_{\parallel\parallel\perp}, \\
\mathcal{L}_n^2 Z^{\varphi} - \partial_{\parallel}^2 Z^{\varphi} - \partial_{\perp}^2 Z^{\varphi} &\simeq 0.
\end{aligned}$$

This second-order system can be written in first-order form by introducing new variables for the temporal and spatial derivatives of the fundamental variables. Absorbing boundary conditions can then be imposed in the $\parallel$ direction, which read (after replacing the new variables with their definitions as first derivatives of the fundamental variables)

$$\mathcal{L}_n \theta \doteq -\partial_{\parallel} \theta, \quad (39)$$

$$\mathcal{L}_n Z_{\parallel} \doteq -\partial_{\parallel} (Z_{\parallel} + D_{\perp\perp\parallel}) - \frac{1}{2} \partial_{\perp} (A_{\perp} - D_{\perp\parallel\parallel} + D_{\perp\perp\perp} - 2D_{\parallel\parallel\perp} + L_{\perp}), \quad (40)$$

$$\begin{aligned}
\mathcal{L}_n Z_{\perp} &\doteq -\partial_{\parallel} [Z_{\perp} - \frac{1}{2} (A_{\perp} + D_{\perp\parallel\parallel} - D_{\perp\perp\perp} + L_{\perp})] \\
&\quad - \frac{1}{2} \partial_{\perp} (A_{\parallel} + D_{\parallel\parallel\parallel} - D_{\parallel\perp\perp} - 2D_{\perp\perp\parallel} + L_{\parallel}), \quad (41)
\end{aligned}$$

$$\mathcal{L}_n Z^{\varphi} \doteq -\partial_{\parallel} Z^{\varphi}, \quad (42)$$

where $\doteq$ denotes equality at the boundary.

The straightforward way of setting up stable outer boundary conditions for the *main* evolution system is to set all ingoing modes to zero at the outer boundaries of the computational domain (so-called *absorbing* boundary conditions). For a symmetric hyperbolic system, this leads to a well-posed initial boundary value problem (IBVP) [17, 18]. Our goal is to investigate under which circumstances such boundary conditions *imply* the constraint-preserving first-order conditions (39–42). Suppose l^- is an incoming mode with characteristic speed $-\lambda$ ($\lambda > 0$). Setting l^- to zero at the entire boundary at all times implies that

$$0 \doteq \partial_{\perp} l^- \equiv L_1(l^-) \quad (43)$$

and

$$0 \doteq \mathcal{L}_n l^- \simeq \lambda \partial_{\parallel} l^- - \partial_{\perp} \mathcal{F}^{\perp}_{l^-} \equiv L_2(l^-), \quad (44)$$

where $\doteq$ denotes equality at the boundary and we have used the general evolution equation for l^- . The relations (43) and (44) will now be used to manipulate the general evolution equations for the Z vector at the boundary. As mentioned in section VI, the evolution system decouples into a non-rotational and a rotational part at the level of principal parts, and we consider the two subsystems separately.

Let us start with the general evolution equations for θ and Z_A , decomposed with respect to our boundary-adapted basis:

$$\begin{aligned}
\mathcal{L}_n \theta &\simeq -\partial_{\parallel} (D_{\parallel\perp\perp} - D_{\perp\perp\parallel} + L_{\parallel} - Z_{\parallel}) - \partial_{\perp} (D_{\perp\parallel\parallel} - D_{\parallel\parallel\perp} + L_{\perp} - Z_{\perp}), \\
\mathcal{L}_n Z_{\parallel} &\simeq -\partial_{\parallel} (\chi_{\perp\perp} + K_{\varphi}^{\varphi} - \theta) + \partial_{\perp} \chi_{\parallel\perp}, \\
\mathcal{L}_n Z_{\perp} &\simeq \partial_{\parallel} \chi_{\parallel\perp} - \partial_{\perp} (\chi_{\parallel\parallel} + K_{\varphi}^{\varphi} - \theta).
\end{aligned}$$

If we add to these suitable combinations of equations (43) and (44),

$$\begin{aligned}
\mathcal{L}_n \theta &+ = L_2(-l_{1,1}^-), \\
\mathcal{L}_n Z_{\parallel} &+ = L_2(-l_{1,1}^- + l_{1,3}^- + l_{1,4}^-) + L_1(-l_{1,2}^-), \\
\mathcal{L}_n Z_{\perp} &+ = L_2(-l_{1,2}^-) + L_1(l_{1,1}^- - l_{1,3}^- - l_f^-),
\end{aligned}$$

we obtain precisely the constraint-preserving boundary conditions (39–41), *provided that we choose harmonic gauge* $f = 1$, $m = 2$, $\frac{m-2}{f-1} = 0$. (For different values of f and m , the derivatives tangential to the boundary in the evolution

equation for $Z_\perp$ differ from the constraint-preserving boundary condition (41).) Hence absorbing boundary conditions for the non-rotational part of the evolution system preserve the constraints.

However, this is not true for the rotational subsystem. A little calculation shows that the only dissipative boundary condition consistent with (42) is

$$l_{1,5}^- \doteq 0, \quad l_{1,6}^- \doteq l_{1,6}^+ \Leftrightarrow B^\varphi \doteq 0.$$

Fortunately, there is a much simpler solution to this problem: we delete Z^φ entirely from the evolution system. Note that setting $Z^\varphi \equiv 0$ does not break general covariance because our spacetime has a Killing vector and we simply choose $Z_\alpha \xi^\alpha = 0$. The remaining rotational subsystem is still symmetric hyperbolic, and the constraint

$$\mathcal{C}_\varphi \simeq \frac{1}{2} E^A{}_{,A}$$

has zero speed with respect to the boundary:

$$\mathcal{L}_n \mathcal{C}_\varphi \simeq \frac{1}{2} \epsilon^{AB} B^\varphi{}_{,AB} = 0.$$

The absorbing boundary condition

$$l_{1,6}^- = E^\perp + B^\varphi \doteq 0 \tag{45}$$

leads to a well-posed IBVP for this system.

In addition to the Z constraints, there are differential constraints related to the definition of the first-order variables,

$$\begin{aligned} D_{ABC} &= \frac{1}{2} \partial_A H_{BC}, \\ L_A &= \lambda^{-1} \partial_A \lambda, \\ A_A &= \alpha^{-1} \partial_A \alpha, \end{aligned}$$

and the ordering constraints (38). By virtue of the evolution equations (10, 11, 26) and (27, 28, 29), it is easy to see that the differential and ordering constraints have zero speed with respect to the boundary.

In summary, we have shown that for harmonic gauge, absorbing boundary conditions for the non-rotational part of the main evolution system are automatically constraint-preserving, and that there are two ways of setting up constraint-preserving stable outer boundary conditions for the rotational subsystem. We thus have a well-posed IBVP with constraint-preserving boundary conditions. The result concerning the non-rotational part carries over immediately to the general Z4 system without spacetime symmetries. Constraint-preserving boundary conditions for the Z4 system have recently also been analysed by Bona et al. [4]: in essence, those authors propose to implement the constraint-preserving boundary conditions (39–41) directly. While it is not clear analytically that such boundary conditions lead to a well-posed IBVP, Bona et al. are able to prove a necessary condition and to demonstrate numerical stability.

VIII. REGULARITY ON AXIS

Extra care is needed in axisymmetric situations because the coordinate system that is adapted to the symmetry, cylindrical polar coordinates, is singular on the axis. This leads to rather strong regularity conditions on tensor fields, as explained in appendix A. In addition to the axisymmetry, we would like to impose reflection symmetry about the z -axis so that we only need to evolve one half of the (r, z) plane. This implies that tensor components are either odd or even functions of z .

Let us first deal with one of the regularity conditions for 2-tensors $M_{\alpha\beta}$, which follows from (1)

$$\frac{M_{\varphi\varphi}}{r^2 M_{rr}} = 1 + O(r^2)$$

near the r -axis. For the metric $g_{\alpha\beta}$ this implies

$$\frac{\lambda^2}{r^2 H_{rr}} = \frac{g_{\varphi\varphi}}{r^2 g_{rr}} = 1 + O(r^2).$$

We enforce this condition by replacing λ with a new variable $\tilde{s}$ defined by

$$\lambda = r e^{r^2 \tilde{s}} \sqrt{H_{rr}}. \tag{46}$$

To satisfy the corresponding condition for the extrinsic curvature, we introduce $\tilde{Y}$ via

$$K_\varphi^\varphi = \frac{\chi_{rr}}{H_{rr}} + r^2 \tilde{Y} \quad (47)$$

(note that $K_{\varphi\varphi} = \lambda^2 K_\varphi^\varphi$). Similarly for the energy-momentum tensor, we set

$$\tau = \frac{S_{rr}}{H_{rr}} + r^2 \tilde{\tau}.$$

We remark that the definitions of the variables $\tilde{s}$ (46) and $\tilde{Y}$ (47) can be viewed as a generalization of those in [5] and [10].

The remaining dependent variables are redefined by taking out systematically the leading order of r and z :

$$\begin{aligned} \tilde{H}_{rr} &= H_{rr}, & \tilde{H}_{rz} &= H_{rz}/(rz), & \tilde{H}_{zz} &= H_{zz}, \\ \tilde{D}_{rrr} &= \frac{1}{2} \tilde{H}_{rr,r}/r, & \tilde{D}_{rrz} &= \frac{1}{2} \tilde{H}_{rz,r}/r, & \tilde{D}_{rzz} &= \frac{1}{2} \tilde{H}_{zz,r}/r, \\ \tilde{D}_{zrr} &= \frac{1}{2} \tilde{H}_{rr,z}/z, & \tilde{D}_{zrz} &= \frac{1}{2} \tilde{H}_{rz,z}/z, & \tilde{D}_{zzz} &= \frac{1}{2} \tilde{H}_{zz,z}/z, \\ \tilde{s}_r &= \tilde{s}_{,r}/r, & \tilde{s}_z &= \tilde{s}_{,z}/z, \\ \tilde{\alpha} &= \alpha, & \tilde{A}_r &= \tilde{\alpha}^{-1} \tilde{\alpha}_{,r}/r, & \tilde{A}_z &= \tilde{\alpha}^{-1} \tilde{\alpha}_{,z}/z, \\ \tilde{\chi}_{rr} &= \chi_{rr}, & \tilde{\chi}_{rz} &= \chi_{rz}/(rz), & \tilde{\chi}_{zz} &= \chi_{zz}, \\ \tilde{E}^r &= E^r/r, & \tilde{E}^z &= E^z/z, & \tilde{B}^\varphi &= B^\varphi/(rz), \\ \tilde{\theta} &= \theta, & \tilde{Z}_r &= Z_r/r, & \tilde{Z}_z &= Z_z/z, & \tilde{Z}^\varphi &= Z^\varphi, \\ \tilde{\sigma} &= \sigma, & \tilde{\rho}_K &= \rho_H - \sigma, & \tilde{J}_r &= J_r/r, & \tilde{J}_z &= J_z/z, & \tilde{J}^\varphi &= J^\varphi, \\ \tilde{\Sigma}_r &= \Sigma_r/r, & \tilde{\Sigma}_z &= \Sigma_z/z, & \tilde{S}_r &= S_r/r, & \tilde{S}_z &= S_z/z, \\ \tilde{S}_{rr} &= S_{rr}, & \tilde{S}_{rz} &= S_{rz}/(rz), & \tilde{S}_{zz} &= S_{zz}. \end{aligned}$$

It can now be verified⁴ that the Z(2+1)+1 equations can be written in terms of the new variables $\tilde{\mathbf{u}}$ in the form

$$\partial_t \tilde{\mathbf{u}} + \left[\alpha \tilde{\mathcal{F}}^{(r^2)}(\tilde{\mathbf{u}}) \right]_{,r^2} + \left[\alpha \tilde{\mathcal{F}}^{(z^2)}(\tilde{\mathbf{u}}) \right]_{,z^2} = \alpha \tilde{\mathcal{S}}(\tilde{\mathbf{u}}), \quad (48)$$

where the fluxes $\tilde{\mathcal{F}}$ and the source $\tilde{\mathcal{S}}$ are even in r and z and manifestly regular on the axes (cf. appendix C).

The above transformation of variables does not affect the hyperbolicity of the system. To see this one should note that

$$\begin{aligned} \tilde{D}_{rrr} &\simeq D_{rrr}/r, & \tilde{D}_{rrz} &\simeq D_{rrz}/(r^2 z), & \tilde{D}_{rzz} &\simeq D_{rzz}/r, \\ \tilde{D}_{zrr} &\simeq D_{zrr}/z, & \tilde{D}_{zrz} &\simeq D_{zrz}/(r z^2), & \tilde{D}_{zzz} &\simeq D_{zzz}/z, \\ \tilde{s}_r &\simeq \left(L_r - \frac{D_{rrr}}{H_{rr}} \right) / r^3, & \tilde{s}_z &\simeq \left(L_z - \frac{D_{zrr}}{H_{rr}} \right) / (r^2 z), \\ \tilde{A}_r &\simeq A_r/r, & \tilde{A}_z &\simeq A_z/z, \\ \tilde{\chi}_{rr} &\simeq \chi_{rr}, & \tilde{\chi}_{rz} &\simeq \chi_{rz}/(rz), & \tilde{\chi}_{zz} &\simeq \chi_{zz}, & \tilde{Y} &\simeq (K_\varphi^\varphi - \frac{\chi_{rr}}{H_{rr}}) / r^2, \\ \tilde{E}^r &\simeq E^r/r, & \tilde{E}^z &\simeq E^z/z, & \tilde{B}^\varphi &\simeq B^\varphi/(rz), \\ \tilde{\theta} &\simeq \theta, & \tilde{Z}_r &\simeq Z_r/r, & \tilde{Z}_z &\simeq Z_z/z, & \tilde{Z}^\varphi &\simeq Z^\varphi, \\ \tilde{J}_r &\simeq J_r/r, & \tilde{J}_z &\simeq J_z/z, \\ \tilde{\Sigma}_r &\simeq \Sigma_r/r, & \tilde{\Sigma}_z &\simeq \Sigma_z/z, & \tilde{S}_r &\simeq S_r/r, & \tilde{S}_z &\simeq S_z/z, \\ \tilde{S}_{rr} &\simeq S_{rr}, & \tilde{S}_{rz} &\simeq S_{rz}/(rz), & \tilde{S}_{zz} &\simeq S_{zz}, \end{aligned}$$

where $\simeq$ denotes equality to leading derivative order (the lower-order terms are absorbed in the source terms)⁵. Hence the variables occurring in the fluxes undergo a *linear* transformation, which does not affect the hyperbolicity and leaves the characteristic variables unchanged.

⁴ The calculations are rather lengthy. We used the computer algebra system REDUCE.

⁵ Note that H_{rr} has zero flux and is thus to be treated as a background quantity.

By writing the equations in terms of derivatives with respect to r^2 and z^2 instead of r and z , however, we have ultimately to multiply the fluxes by $2r$ and $2z$, respectively, because

$$\partial_{r^2} = \frac{1}{2r} \partial_r$$

and similarly for z . This means that in (r^2, z^2) coordinates, the characteristic speeds behave like $\sim r$ and $\sim z$, respectively. For computational purposes it is highly desirable to avoid the non-uniform characteristic speeds which would occur on a (r^2, z^2) grid. We therefore recommend working in the original (r, z) grid and discretizing the derivatives in (48) with respect to r^2 and z^2 . This has another, perhaps more fundamental, advantage: in (r, z) coordinates, we can enforce Neumann boundary conditions for all our variables on the axes (since they are even functions of r and z), whereas in (r^2, z^2) coordinates the boundary conditions on the axes are not known. The authors of [15] faced a similar problem but decided to work on an (r^2, z^2) grid, using extrapolation on the axis, which led to numerical instabilities.

Choptuik et al [5] and Garfinkle et al [10] take a slightly different approach. They include terms which are formally irregular on axis, but which are regular once appropriate boundary conditions are imposed. Following an idea of Evans [7] these terms are differenced with respect to r^2 or r^3 .

The transformation from conserved to primitive fluid variables and vice versa can be made manifestly regular on the axes by redefining the velocities

$$\tilde{v}_r = v_r/r, \quad \tilde{v}_z = v_z/z$$

and leaving the remaining primitive variables unchanged. Note in particular that the modified general matter variable

$$\tilde{\tau} = r^{-2} \left(\tau - \frac{S_{rr}}{H_{rr}} \right) = \rho h W^2 \left(e^{2r^2 \tilde{s}} H_{rr} v^{\varphi^2} - \frac{\tilde{v}_r^2}{H_{rr}} \right) \quad (49)$$

is then automatically regular.

To compute the characteristic variables (geometry and matter), one can start from the regularized conserved variables $\tilde{\mathbf{u}}$, compute the original conserved variables $\mathbf{u}$ and evaluate the characteristic variables (section VI and appendix D 3). While this transformation is perfectly regular on the axes, the inverse transformation involves negative powers of r and z . In order to implement outer boundary conditions at $r = r_{max}$ ($z = z_{max}$), one has to ensure that the inverse transformation in the r -direction (z -direction) is regular at $z = 0$ ($r = 0$). This can be achieved by redefining the *characteristic* variables in a similar fashion as was done for the conserved variables above. We choose to make the following replacements

- for the characteristic variables in the r -direction:

$$l_{0,4} \rightarrow \tilde{l}_{0,4} = z^{-2} \left(l_{0,4} + \frac{z \tilde{H}_{rz}}{\sqrt{H}} l_{0,5} \right),$$

and the leading order of z is taken out of the remaining characteristic variables

- for the characteristic variables in the z -direction:

$$l_{0,4} \rightarrow \tilde{l}_{0,4} = r^{-2} \left(l_{0,4} + \frac{r \tilde{H}_{rz}}{\sqrt{H}} l_{0,5} \right),$$

$$l_{0,6} \rightarrow \tilde{l}_{0,6} = r^{-3} (l_{0,6} - l_{0,5}),$$

$$l_{1,4}^{\pm} \rightarrow \tilde{l}_{1,4}^{\pm} = r^{-2} (l_{1,4}^{\pm} - l_{1,3}^{\pm}),$$

and the leading order of r is taken out of the remaining characteristic variables

We have checked with REDUCE that the transformation from characteristic variables to conserved variables and *vice versa* is then manifestly regular at the entire outer boundary.

However, the transformation from characteristic variables in the r -direction (z -direction) to conserved variables is still formally singular at $r = 0$ ($z = 0$), and no regularization procedure can cure this problem. To understand this, one can note that the characteristic variables in the r -direction (z -direction) do not have a definite parity in r (z). For this reason, numerical methods operating in the space of characteristic variables (typically ones based on the solution of the Riemann problem) appear to be unusable near the axes.

IX. CONCLUSIONS

We started out by trying to understand why so many previous attempts to evolve axisymmetric spacetimes failed because of on-axis instabilities. This led to a detailed study of the behaviour of the components of axisymmetric tensors with respect to cylindrical polar coordinates, given that the components with respect to cartesian coordinates were regular in the neighbourhood of the axis. This suggested, most strongly, that we should make a Kaluza-Klein-like reduction to a 3-dimensional Lorentzian spacetime [11], a step taken earlier by Maeda et al [14], [15]. (A different version of this idea, without rotation, has been pursued by [5].)

Choptuik et al's evolution [5] was fully constrained—elliptic equations were solved for the metric components, making use of multigrid for computational efficiency. We made similar choices and obtained similar results. However in strong field situations, e.g., near-critical collapse of Brill waves, the diagonal dominance of the matrix describing the discretized elliptic operators failed, and so the multigrid iterations failed to converge.

We have not found a satisfactory resolution of this problem, and so we went to the other extreme—our evolution algorithm involves no elliptic operators. To achieve this we have used the Z4 formalism [2], [3] adapted to our axisymmetric reduction, an evolution equation for the lapse function and a zero shift gauge. Our first order hyperbolic system of conservation laws (with sources) is strongly hyperbolic and, for one choice of parameters, symmetric hyperbolic. In addition the dependent variables have been carefully chosen so that each and every term in the system is regular on axis. We have allowed for arbitrary matter sources and have presented explicitly the details for a perfect fluid.

Initial conditions have to be imposed via the constraint conditions, which means solving elliptic equations. By means of conformal rescalings on the initial hypersurface one can avoid the diagonal dominance problem on that hypersurface and so multigrid techniques are applicable.

One must choose a high resolution shock capturing method for the evolution of the hyperbolic system. Our system contains one subtle disadvantage if we choose to work with numerical algorithms that require a transformation between conserved and characteristic variables. The conversion from characteristic variables to our “regularized” conserved variables, see section VIII, is formally singular on the axis. However, we have shown that the transformation can be used to impose outer boundary conditions in a regular way.

Because the evolution does not involve elliptic operators it is straightforward to introduce adaptive mesh refinement, where appropriate, to enhance the resolution.

We shall be using our formalism to evolve a number of interesting axisymmetric spacetimes, including the effects of rotation and perfect fluid sources.

APPENDIX A: CONSEQUENCES OF AXISYMMETRY

We want to use a (t, z, r, φ) chart adapted to the Killing vector $\xi = \partial/\partial\varphi$. Unfortunately this chart is singular at the axis $r = 0$. We shall assume *elementary flatness*: in a neighbourhood of the axis we can introduce local cartesian coordinates $x^A = (x, y)$ such that

$$x = r \cos \varphi, \quad y = r \sin \varphi \quad \Longleftrightarrow \quad r = \sqrt{x^2 + y^2}, \quad \varphi = \arctan \frac{y}{x}.$$

With respect to cartesian coordinates the Killing vector is

$$\xi = -y\partial/\partial x + x\partial/\partial y.$$

Notice that this representation is valid everywhere, while $\xi = \partial/\partial\varphi$ is valid only for $r > 0$.

We say that a scalar function $f(x^A)$ is *regular on axis* if f has a Taylor expansion with respect to x and y about $x^A = 0$ convergent in some neighbourhood of $r = 0$. (Throughout this section we are ignoring t, z dependencies which are implicit in all calculations.) If f is axisymmetric

$$\mathcal{L}_\xi f = 0 = \partial f / \partial \varphi \quad \Rightarrow \quad f = f(r), \quad r > 0. \quad (\text{A1})$$

If f had a Taylor expansion in r and was regular on axis then the expansion could contain only even powers of r since $r = \sqrt{x^2 + y^2}$, and r has no such expansion. While the conclusion is correct, the argument is fallacious because equation (A1) is invalid at $r = 0$. Instead we start from

$$K \equiv -y f_{,x} + x f_{,y} = 0, \quad (\text{A2})$$

which is valid everywhere. In particular we may differentiate (A2) an arbitrary number of times with respect to x and also with respect to y . Then setting $x^A = 0$ we obtain linear recurrence relations between the Taylor coefficients on axis. These can be solved to show $f = \sum_n f_n (x^2 + y^2)^n$.

Next consider a vector field u^α . For $a = (t, z)$, $\mathcal{L}_\xi u^a = 0$ implies $\partial u^a / \partial \varphi = 0$. This reduces to the scalar field case and we may deduce $u^a = u^a(r^2)$. For u^x and u^y we have

$$\partial u^x / \partial \varphi + u^y = 0, \quad \partial u^y / \partial \varphi - u^x = 0.$$

The general solution for $r > 0$ is

$$u^x = \hat{a}(r) \cos \varphi - \hat{b}(r) \sin \varphi, \quad u^y = \hat{a}(r) \sin \varphi + \hat{b}(r) \cos \varphi.$$

However in the cartesian chart we have

$$-y u^x_{,x} + x u^x_{,y} + u^y = 0, \quad -y u^y_{,x} + x u^y_{,y} - u^x = 0. \quad (\text{A3})$$

Clearly $u^A = 0$ on axis. We write $\hat{a} = ra$ etc so that

$$u^x = xa - yb, \quad u^y = ya + xb. \quad (\text{A4})$$

We now regard a and b as unknown functions of x and y to be determined by substituting (A4) into (A3), differentiating the latter an arbitrary number of times, and then solving the recurrence relations for the Taylor coefficients of a and b . Again we find that a and b are even functions of r . Thus in the (t, z, r, φ) chart an axisymmetric vector field which is regular on axis must take the form

$$u^\alpha = (A, B, rC, D), \quad (\text{A5})$$

where A, B, C and D are functions of t, z and r^2 .

Next consider an axisymmetric covector field ω_α which is regular on axis. For $a = (t, z)$, $\mathcal{L}_\xi \omega_a = 0$ implies $\partial \omega_a / \partial \varphi = 0$. This reduces to the scalar field case and we may deduce $\omega_a = \omega_a(r^2)$. For the other indices we find

$$-y \omega_{x,x} + x \omega_{x,y} + \omega_y = 0, \quad -y \omega_{y,x} + x \omega_{y,y} - \omega_x = 0, \quad (\text{A6})$$

which is equivalent to (A3), interchanging u^A and ω_A . We therefore deduce the analogue of (A4) and hence, in polar coordinates,

$$\omega_\alpha = (E, F, rG, r^2H), \quad (\text{A7})$$

where E , F , G and H are functions of t , z and r^2 .

Finally we consider a symmetric valence 2 axisymmetric tensor field $M_{\alpha\beta}$ which is both axisymmetric and regular on axis. For $(a, b) = (t, z)$ we have $\mathcal{L}_\xi M_{ab} = 0$ and so $M_{ab} = M_{ab}(r^2)$. Letting $a = (t, z)$ we have

$$-yM_{ax,x} + xM_{ax,y} + M_{ay} = 0, \quad -yM_{ay,x} + xM_{ay,y} - M_{ax} = 0. \quad (\text{A8})$$

This is essentially the same as (A6) and we may deduce $M_{ar} = rA_a(r^2)$ and $M_{a\varphi} = r^2B_a(r^2)$. The remaining Killing equations are

$$-yM_{xx,x} + xM_{xx,y} + 2M_{xy} = 0, \quad -yM_{yy,x} + xM_{yy,y} - 2M_{xy} = 0, \quad -yM_{xy,x} + xM_{xy,y} + M_{yy} - M_{xx} = 0.$$

If we introduce new variables $u = \frac{1}{2}(M_{xx} + M_{yy})$, $v = \frac{1}{2}(M_{xx} - M_{yy})$ and $w = M_{xy}$ then

$$-yu_{,x} + xu_{,y} = 0$$

which implies $u = u(r^2)$. The remaining equations are

$$-yv_{,x} + xv_{,y} + 2w = 0, \quad -yw_{,x} + xw_{,y} - 2v = 0. \quad (\text{A9})$$

For $r > 0$ these can be written as

$$v_{,\varphi} + 2w = 0, \quad w_{,\varphi} - 2v = 0,$$

so that

$$v = \hat{a}(r) \cos 2\varphi - \hat{b}(r) \sin 2\varphi, \quad w = \hat{a}(r) \sin 2\varphi + \hat{b}(r) \cos 2\varphi,$$

where $\hat{a}$ and $\hat{b}$ are arbitrary functions of r . If we set $a = \hat{a}/r^2$ and $b = \hat{b}/r^2$ then

$$v = (x^2 - y^2)a - 2xyb, \quad w = 2xya + (x^2 - y^2)b. \quad (\text{A10})$$

(Note that (A9) and its first derivatives imply that v , w and their first derivatives vanish on axis, consistent with (A10).) Substituting (A10) into (A9) gives

$$x^3a_{,y} - x^2y(a_{,x} + 2b_{,y}) - xy^2(a_{,y} - 2b_{,x}) + y^3a_{,x} = 0, \quad (\text{A11})$$

$$x^3b_{,y} + x^2y(-b_{,x} + 2a_{,y}) + xy^2(-2a_{,x} - b_{,y}) + y^3b_{,x} = 0. \quad (\text{A12})$$

Differentiating these many times and, proceeding as in the scalar and vector cases we conclude that a and b are functions of r^2 . Thus

$$M_{xx} = u + (x^2 - y^2)a - 2xyb, \quad M_{yy} = u - (x^2 - y^2)a + 2xyb, \quad M_{xy} = 2xya + (x^2 - y^2)b.$$

Re-expressing these as polar components we obtain

$$M_{rr} = u + r^2a, \quad M_{r\varphi} = r^3b, \quad M_{\varphi\varphi} = r^2(u - r^2a). \quad (\text{A13})$$

Finally combining all of the results we have

$$M_{\alpha\beta} = \begin{pmatrix} A & B & rD & r^2F \\ B & C & rE & r^2G \\ rD & rE & H + r^2J & r^3K \\ r^2F & r^2G & r^3K & r^2(H - r^2J) \end{pmatrix}, \quad (\text{A14})$$

where $A, B, \dots, K$ are functions of t , z and r^2 .

APPENDIX B: TRANSFORMATION FROM CHARACTERISTIC TO CONSERVED VARIABLES

The conserved variables are given in terms of the characteristic variables by

$$\begin{aligned}
D_{\parallel\parallel\parallel} &= -\frac{1}{f}l_{0,1} + \left(\frac{f(m-2)}{2(f-1)} + 1\right)(l_{1,1}^+ - l_{1,1}^-) - \frac{1}{2}(l_{1,3}^+ - l_{1,3}^- + l_{1,4}^+ - l_{1,4}^-) \\
&\quad + \frac{1}{2f}(l_f^+ + l_f^-), \\
D_{\parallel\parallel\perp} &= -\frac{1}{fm}l_{0,2} + \frac{(m-2)}{2m}(l_{0,3} + l_{0,5} + l_{0,6}) - \frac{(fm-2)}{2fm}l_{0,7} + \frac{1}{2}(l_{1,2}^+ - l_{1,2}^-), \\
D_{\parallel\perp\perp} &= \frac{1}{2}(l_{1,3}^+ - l_{1,3}^-), \\
D_{\perp\parallel\parallel} &= l_{0,3}, \\
D_{\perp\perp\parallel} &= l_{0,4}, \\
D_{\perp\perp\perp} &= l_{0,5}, \\
L_{\parallel} &= \frac{1}{2}(l_{1,4}^+ - l_{1,4}^-), \\
L_{\perp} &= l_{0,6}, \\
A_{\parallel} &= \frac{f(m-2)}{2(f-1)}(l_{1,1}^+ - l_{1,1}^-) + \frac{1}{2}(l_f^+ + l_f^-), \\
A_{\perp} &= l_{0,7}, \\
\chi_{\parallel\parallel} &= \left(\frac{f(m-2)}{2(f-1)} + 1\right)(l_{1,1}^+ + l_{1,1}^-) - \frac{1}{2}(l_{1,3}^+ + l_{1,3}^- + l_{1,4}^+ + l_{1,4}^-) \\
&\quad + \frac{1}{2\sqrt{f}}(l_f^+ - l_f^-), \\
\chi_{\parallel\perp} &= \frac{1}{2}(l_{1,2}^+ + l_{1,2}^-), \\
\chi_{\perp\perp} &= \frac{1}{2}(l_{1,3}^+ + l_{1,3}^-), \\
K_{\varphi}^{\varphi} &= \frac{1}{2}(l_{1,4}^+ + l_{1,4}^-), \\
E^{\parallel} &= \frac{1}{2}(l_{1,5}^+ + l_{1,5}^-), \\
E^{\perp} &= \frac{1}{2}(l_{1,6}^+ + l_{1,6}^-), \\
B^{\varphi} &= -\frac{1}{2}(l_{1,6}^+ - l_{1,6}^-), \\
\theta &= \frac{1}{2}(l_{1,1}^+ + l_{1,1}^-), \\
Z_{\parallel} &= -l_{0,4} + \frac{1}{2}(-l_{1,1}^+ + l_{1,1}^- + l_{1,3}^+ - l_{1,3}^- + l_{1,4}^+ - l_{1,4}^-), \\
Z_{\perp} &= \frac{1}{2}(l_{0,3} - l_{0,5} + l_{0,6} + l_{0,7}) - \frac{1}{2}(l_{1,2}^+ - l_{1,2}^-), \\
Z^{\varphi} &= -\frac{1}{4}(l_{1,5}^+ - l_{1,5}^-).
\end{aligned}$$

Tensor components can then be computed as, for instance,

$$\chi_{AB} = \mu_A \mu_B \chi_{\parallel\parallel} + 2\mu_{(A} \pi_{B)} \chi_{\parallel\perp} + \pi_A \pi_B \chi_{\perp\perp}.$$

APPENDIX C: REGULAR CONSERVATION FORM

In this appendix we provide some details of the regular form of the Z(2+1)+1 equations outlined in section VIII. The equations are written in conservation form (48)

$$\partial_t \tilde{\mathbf{u}} + \left[\alpha \tilde{\mathcal{F}}^{(r^2)}(\tilde{\mathbf{u}}) \right]_{,r^2} + \left[\alpha \tilde{\mathcal{F}}^{(z^2)}(\tilde{\mathbf{u}}) \right]_{,z^2} = \alpha \tilde{\mathcal{S}}(\tilde{\mathbf{u}}),$$

where the modified variables $\tilde{\mathbf{u}}$ are defined in section VIII.

The fluxes $\tilde{\mathcal{F}}^{(r^2)}$, $\tilde{\mathcal{F}}^{(z^2)}$ and sources $\tilde{\mathcal{S}}$ have been obtained with the computer algebra system REDUCE and have been typeset automatically using the TeX REDUCE Interface (TRI). From the same source we generated C++ code

using the REDUCE source code optimization packages SCOPE and GENTRAN. The resulting expressions are rather lengthy. To get some idea of what is typically involved we state here the fluxes and source for the variable $\tilde{Y}$.

Let us introduce the shorthand

$$H = \tilde{H}_{rr}\tilde{H}_{zz} - r^2 z^2 \tilde{H}_{rz}^2$$

for the determinant of the 2-metric. The fluxes are given by

$$\begin{aligned}\tilde{\mathcal{F}}_{\tilde{Y}}^{(r^2)} &= 2H^{-1}\tilde{D}_{rrr}r^2z^2\tilde{H}_{rr}^{-2}\tilde{H}_{rz}^2 - 2H^{-1}\tilde{D}_{rzz} - 4H^{-1}\tilde{D}_{zrr}z^2\tilde{H}_{rr}^{-1}\tilde{H}_{rz} + 4H^{-1}\tilde{D}_{zrz}z^2 \\ &\quad + 2\tilde{H}_{rr}^{-1}(-\tilde{A}_r + 2\tilde{Z}_r) + 2H^{-1}r^2z^2\tilde{H}_{rz}(r^2\tilde{H}_{rr}^{-1}\tilde{H}_{rz}\tilde{s}_r - \tilde{s}_z), \\ \tilde{\mathcal{F}}_{\tilde{Y}}^{(z^2)} &= 2H^{-1}z^2(-r^2\tilde{H}_{rz}\tilde{s}_r + \tilde{H}_{rr}\tilde{s}_z),\end{aligned}$$

and the source term is

$$\begin{aligned}\tilde{\mathcal{S}}_{\tilde{Y}} &= -\kappa\tilde{\tau} + 2H^{-1}\tilde{\chi}_{rr}z^2\tilde{H}_{rr}^{-2}\tilde{H}_{rz}^2 + 2\tilde{\chi}_{rr}\tilde{H}_{rr}^{-1}\tilde{Y} + H^{-1}\tilde{\chi}_{rr}z^2\tilde{H}_{rr}^{-1}\tilde{H}_{rz}(r^2\tilde{H}_{rz}\tilde{Y} - 4\tilde{\chi}_{rz}) \\ &\quad + H^{-1}\tilde{\chi}_{zz}\tilde{H}_{rr}\tilde{Y} - 4H^{-1}\tilde{D}_{rrr}r^2z^2\tilde{H}_{rr}^{-3}\tilde{H}_{rz}^2 \\ &\quad - H^{-2}\tilde{D}_{rrr}^2r^4z^4\tilde{H}_{rr}^{-3}\tilde{H}_{rz}^4 + 4H^{-1}\tilde{D}_{rrr}\tilde{D}_{rrz}r^2z^2\tilde{H}_{rr}^{-2}\tilde{H}_{rz} \\ &\quad + 2H^{-2}\tilde{D}_{rrr}\tilde{D}_{rrz}r^4z^4\tilde{H}_{rr}^{-2}\tilde{H}_{rz}^3 + 4H^{-1}\tilde{D}_{rrr}\tilde{D}_{zrr}z^2\tilde{H}_{rr}^{-2}\tilde{H}_{rz} \\ &\quad + 2H^{-2}\tilde{D}_{rrr}\tilde{D}_{zrr}r^2z^4\tilde{H}_{rr}^{-2}\tilde{H}_{rz}^3 - 2H^{-2}\tilde{D}_{rrr}\tilde{D}_{zzz}z^2\tilde{H}_{rz} \\ &\quad + H^{-1}\tilde{D}_{rrr}\tilde{H}_{rr}^{-2}\tilde{H}_{rz}(-5r^4z^2\tilde{H}_{rz}\tilde{s}_r - r^2z^2\tilde{A}_r\tilde{H}_{rz} + 4r^2z^2\tilde{H}_{rr}\tilde{s}_z - 2r^2z^2\tilde{H}_{rz}\tilde{s} \\ &\quad \quad + 4r^2z^2\tilde{H}_{rz}\tilde{Z}_r + 2z^2\tilde{A}_z\tilde{H}_{rr} - 4z^2\tilde{H}_{rr}\tilde{Z}_z + 2\tilde{H}_{rr}) \\ &\quad + H^{-2}\tilde{D}_{rrr}r^2z^2\tilde{H}_{rr}^{-2}\tilde{H}_{rz}^3(-r^4z^2\tilde{H}_{rz}\tilde{s}_r + r^2z^2\tilde{H}_{rr}\tilde{s}_z + 2r^2z^2\tilde{H}_{rz}\tilde{s} + 2z^2\tilde{H}_{rz} + 2\tilde{H}_{rr}) \\ &\quad - 2H^{-2}\tilde{D}_{rrz}\tilde{D}_{rzz}r^2z^2\tilde{H}_{rz} - 4H^{-1}\tilde{D}_{rrz}\tilde{D}_{zrr}z^2\tilde{H}_{rr}^{-1} - 6H^{-2}\tilde{D}_{rrz}\tilde{D}_{zrr}r^2z^4\tilde{H}_{rr}^{-1}\tilde{H}_{rz}^2 \\ &\quad + 4H^{-2}\tilde{D}_{rrz}\tilde{D}_{zrz}r^2z^4\tilde{H}_{rz} + 2H^{-2}\tilde{D}_{rrz}\tilde{D}_{zzz}z^2\tilde{H}_{rr} \\ &\quad + 2H^{-1}\tilde{D}_{rrz}(2r^4z^2\tilde{H}_{rr}^{-1}\tilde{H}_{rz}\tilde{s}_r + r^2z^2\tilde{A}_r\tilde{H}_{rr}^{-1}\tilde{H}_{rz} - 2r^2z^2\tilde{H}_{rr}^{-1}\tilde{H}_{rz}\tilde{Z}_r \\ &\quad \quad - r^2z^2\tilde{s}_z - z^2\tilde{A}_z + 2z^2\tilde{Z}_z - 1) \\ &\quad + 2H^{-2}\tilde{D}_{rrz}r^2z^2\tilde{H}_{rz}^2(r^4z^2\tilde{H}_{rr}^{-1}\tilde{H}_{rz}\tilde{s}_r - 2r^2z^2\tilde{H}_{rr}^{-1}\tilde{H}_{rz}\tilde{s} - r^2z^2\tilde{s}_z - z^2\tilde{H}_{rr}^{-1}\tilde{H}_{rz} - 1) \\ &\quad + H^{-2}\tilde{D}_{rzz}^2\tilde{H}_{rr} + 4H^{-2}\tilde{D}_{rzz}\tilde{D}_{zrr}z^2\tilde{H}_{rz} - 4H^{-2}\tilde{D}_{rzz}\tilde{D}_{zrz}z^2\tilde{H}_{rr} \\ &\quad + H^{-1}\tilde{D}_{rzz}(-r^2\tilde{s}_r - \tilde{A}_r - 2\tilde{s}) + H^{-2}\tilde{D}_{rzz}r^2z^2\tilde{H}_{rz}(-r^2\tilde{H}_{rz}\tilde{s}_r + \tilde{H}_{rr}\tilde{s}_z + 2\tilde{H}_{rz}\tilde{s}) \\ &\quad + 2H^{-1}\tilde{D}_{zrr}z^2(r^2\tilde{H}_{rr}^{-1}\tilde{H}_{rz}\tilde{s}_r - \tilde{A}_r\tilde{H}_{rr}^{-1}\tilde{H}_{rz} - \tilde{s}_z) \\ &\quad + H^{-2}\tilde{D}_{zrr}z^4\tilde{H}_{rz}^2(r^4\tilde{H}_{rr}^{-1}\tilde{H}_{rz}\tilde{s}_r - 2r^2\tilde{H}_{rr}^{-1}\tilde{H}_{rz}\tilde{s} - r^2\tilde{s}_z - 4\tilde{H}_{rr}^{-1}\tilde{H}_{rz}) \\ &\quad + 2H^{-1}\tilde{D}_{zrz}z^2(\tilde{A}_r + 2\tilde{s}) + 2H^{-2}\tilde{D}_{zrz}z^4\tilde{H}_{rz}(-r^4\tilde{H}_{rz}\tilde{s}_r + r^2\tilde{H}_{rr}\tilde{s}_z + 2r^2\tilde{H}_{rz}\tilde{s} + 2\tilde{H}_{rz}) \\ &\quad + H^{-2}\tilde{D}_{zzz}z^2\tilde{H}_{rr}(r^2\tilde{H}_{rz}\tilde{s}_r - \tilde{H}_{rr}\tilde{s}_z - 2\tilde{H}_{rz}\tilde{s}) - \frac{1}{2}e^{2r^2\tilde{s}}Hz^2\tilde{E}z^2 \\ &\quad + e^{2r^2\tilde{s}}r^2(-z^4\tilde{E}z^2\tilde{H}_{rz}^2 + z^2\tilde{B}^{\varphi^2}\tilde{H}_{rr} - 2z^2\tilde{E}^r\tilde{E}^z\tilde{H}_{rr}\tilde{H}_{rz} - \tilde{E}^{r^2}\tilde{H}_{rr}^2) - r^2\tilde{A}_r\tilde{H}_{rr}^{-1}\tilde{s}_r \\ &\quad + 2r^2\tilde{H}_{rr}^{-1}\tilde{s}_r\tilde{Z}_r + r^2\tilde{Y}^2 - 2\tilde{A}_r\tilde{H}_{rr}^{-1}\tilde{s} + 2\tilde{A}_r\tilde{H}_{rr}^{-1}\tilde{Z}_r + 4\tilde{H}_{rr}^{-1}\tilde{s}\tilde{Z}_r - 2\tilde{\Theta}\tilde{Y} \\ &\quad + H^{-1}(-r^6z^2\tilde{H}_{rr}^{-1}\tilde{H}_{rz}^2\tilde{s}_r^2 - 4r^4z^2\tilde{H}_{rr}^{-1}\tilde{H}_{rz}^2\tilde{s}\tilde{s}_r + 2r^4z^2\tilde{H}_{rr}^{-1}\tilde{H}_{rz}^2\tilde{s}_r\tilde{Z}_r \\ &\quad \quad + 2r^4z^2\tilde{H}_{rz}\tilde{s}_r\tilde{s}_z - 2r^2z^2\tilde{A}_r\tilde{H}_{rr}^{-1}\tilde{H}_{rz}^2\tilde{s} - 2r^2z^2\tilde{\chi}_{rz}\tilde{H}_{rz}\tilde{Y} - r^2z^2\tilde{H}_{rr}\tilde{s}_z^2 \\ &\quad \quad - 4r^2z^2\tilde{H}_{rr}^{-1}\tilde{H}_{rz}^2\tilde{s}^2 + 4r^2z^2\tilde{H}_{rr}^{-1}\tilde{H}_{rz}^2\tilde{s}\tilde{Z}_r - 3r^2z^2\tilde{H}_{rr}^{-1}\tilde{H}_{rz}^2\tilde{s}_r \\ &\quad \quad + 4r^2z^2\tilde{H}_{rz}\tilde{s}\tilde{s}_z - 2r^2z^2\tilde{H}_{rz}\tilde{s}_r\tilde{Z}_z - 2r^2z^2\tilde{H}_{rz}\tilde{s}_z\tilde{Z}_r - r^2\tilde{H}_{rz}\tilde{s}_r \\ &\quad \quad + 2z^2\tilde{A}_z\tilde{H}_{rz}\tilde{s} + 2z^2\tilde{\chi}_{rz}^2 + 2z^2\tilde{H}_{rr}\tilde{s}_z\tilde{Z}_z - 6z^2\tilde{H}_{rr}^{-1}\tilde{H}_{rz}^2\tilde{s} \\ &\quad \quad - 4z^2\tilde{H}_{rz}\tilde{s}\tilde{Z}_z + 4z^2\tilde{H}_{rz}\tilde{s}_z + \tilde{H}_{rr}\tilde{s}_z + 2\tilde{H}_{rz}\tilde{s}) \\ &\quad + H^{-2}z^2\tilde{H}_{rz}^2(r^4z^2\tilde{H}_{rr}^{-1}\tilde{H}_{rz}^2\tilde{s}_r - r^4\tilde{H}_{rz}\tilde{s}_r - 2r^2z^2\tilde{H}_{rr}^{-1}\tilde{H}_{rz}^2\tilde{s} - r^2z^2\tilde{H}_{rz}\tilde{s}_z \\ &\quad \quad + r^2\tilde{H}_{rr}\tilde{s}_z + 2r^2\tilde{H}_{rz}\tilde{s} - z^2\tilde{H}_{rr}^{-1}\tilde{H}_{rz}^2).\end{aligned}$$

Note that the expressions are manifestly regular on the axes and even in r and z .

APPENDIX D: MATTER EVOLUTION EQUATIONS

1. Conservation form

The matter evolution equations (16–19) can clearly be written in conservation form

$$\partial_t \mathbf{u} + [\alpha \mathcal{F}^D(\mathbf{u})]_{,D} = \alpha \mathcal{S}(\mathbf{u}).$$

Following [1], we replace ρ_H with $\rho_K = \rho_H - \sigma$ (kinetic energy) and regard as the set of conserved variables

$$\mathbf{u} = (\rho_K, J_A, J^\varphi, \sigma)^T. \quad (\text{D1})$$

The fluxes are

$$\begin{aligned} \mathcal{F}^D_{\rho_K} &= J^D - \Sigma^D, \\ \mathcal{F}^D_{J_A} &= S_A^D, \\ \mathcal{F}^D_{J^\varphi} &= S^D, \\ \mathcal{F}^D_\sigma &= \Sigma^D, \end{aligned}$$

and the source terms are

$$\begin{aligned} \mathcal{S}_{\rho_K} &= (\Sigma^A - J^A)(D^I_A + L_A) + K_\varphi{}^\varphi(\tau - \sigma) + \chi_{AB} S^{AB} - J^A A_A + \chi \rho_K \\ &\quad + \lambda^2 E^A S_A, \\ \mathcal{S}_{J_A} &= -S_{AB}(A^B + L^B) + J_A(\chi + K_\varphi{}^\varphi) - A_A \rho_H + L_A \tau \\ &\quad + \lambda^2 (E_A J^\varphi + \epsilon_{AB} S^B B^\varphi), \\ \mathcal{S}_{J^\varphi} &= -(D^I_A + 3L_A) S^A + J^\varphi(\chi + 3K_\varphi{}^\varphi), \\ \mathcal{S}_\sigma &= -(D^I_A + L_A) \Sigma^A + \sigma(\chi + K_\varphi{}^\varphi). \end{aligned}$$

2. Perfect fluid

To evaluate the characteristic structure, we need to specify the matter model. Here, we consider a perfect fluid with four-velocity u^α , normalized such that

$$u_\alpha u^\alpha = -1,$$

rest mass density ρ , pressure p and internal energy ϵ . The dependence of the pressure on the density and the internal energy is given by the equation of state

$$p = p(\rho, \epsilon). \quad (\text{D2})$$

With those definitions, the number density is

$$N^\alpha = \rho u^\alpha$$

and the energy-momentum tensor is given by

$$T^{\alpha\beta} = \rho h u^\alpha u^\beta + p g^{\alpha\beta},$$

where h is the specific enthalpy,

$$h = 1 + \epsilon + \frac{p}{\rho}. \quad (\text{D3})$$

The Lorentz factor is defined as

$$W \equiv -u^\alpha n_\alpha. \quad (\text{D4})$$

Observers who are at rest in a slice $\Sigma(t)$ (i.e., who have four-velocity n^α) measure a coordinate velocity

$$v^A = W^{-1} h_\alpha^A u^\alpha,$$

and the angular velocity is

$$v^\varphi = W^{-1} \lambda^{-2} \xi_\alpha u^\alpha.$$

Hence we obtain the familiar relation

$$W = (1 - v^2)^{-1/2}, \quad (\text{D5})$$

where

$$v^2 = v_A v^A + \lambda^2 v^\varphi^2.$$

The variables

$$\mathbf{w} = (v_A, v^\varphi, \rho, p, \epsilon, h, W)^T$$

are called *primitive variables*.⁶ The conserved variables can be expressed in terms of the primitive variables as

$$\begin{aligned} \rho_K &= \rho h W^2 - p - \rho W, \\ J_A &= \rho h W^2 v_A, \\ J^\varphi &= \rho h W^2 v^\varphi, \\ \sigma &= \rho W, \end{aligned} \quad (\text{D6})$$

and the remaining matter variables are

$$\begin{aligned} \tau &= \rho h W^2 \lambda^2 v^\varphi^2 + p, \\ S_A &= \rho h W^2 v^\varphi v_A, \\ S_{AB} &= \rho h W^2 v_A v_B + p H_{AB}, \\ \Sigma_A &= \rho W v_A. \end{aligned} \quad (\text{D7})$$

3. Characteristic decomposition

The characteristic decomposition for 3+1 general relativistic hydrodynamics was first derived by the Valencia group [1]. The application to our (2+1)+1 system is straightforward. Our method differs slightly in that we choose a general orthonormal basis (μ^A, π^A) in two-space as in section VI and project vectors along μ (index $\parallel$) and π (index $\perp$). Following the notation of [9], we introduce a few abbreviations. From the equation of state (D2), we form

$$\chi \equiv \frac{\partial p}{\partial \rho}, \quad \kappa \equiv \frac{\partial p}{\partial \epsilon}, \quad h c_s^2 \equiv \chi + \frac{p}{\rho^2} \kappa,$$

where c_s is known as the *sound speed*. Also set⁷

$$\begin{aligned} \mathcal{K}^{-1} &= 1 - \frac{c_s^2 \rho}{\kappa}, \quad \mathcal{V}^\pm = \frac{v_\parallel - \lambda_s^\pm}{1 - v_\parallel \lambda_s^\pm}, \quad \mathcal{A}^\pm = \frac{1 - v_\parallel^2}{1 - v_\parallel \lambda_s^\pm}, \\ \mathcal{C}^\pm &= v_\parallel - \mathcal{V}^\pm, \quad \xi = 1 - v_\parallel^2, \quad \Delta = h^3 W (1 - \mathcal{K}^{-1}) (\mathcal{C}^+ - \mathcal{C}^-) \xi. \end{aligned}$$

⁶ Note only five of those are independent because of (D2), (D3) and (D5).

⁷ Our definitions of ξ and Δ differ from those in [9] by a factor of λ^2 to ensure regularity (see section VIII). We have defined $\mathcal{K}^{-1}$ instead of $\mathcal{K}$ to allow for the special case of the ultrarelativistic equation of state, for which $\mathcal{K}^{-1} = 0$. As a consequence, Δ above has been multiplied by $\mathcal{K}^{-1}$ and the characteristic variable $l_{0,1}$ has been divided by $\mathcal{K}^{-1}$.

The system is found to be strongly hyperbolic. The characteristic speeds in the μ -direction are

$$\begin{aligned}\lambda_0 &= v_{\parallel}, \\ \lambda_s^{\pm} &= \frac{1}{1 - v^2 c_s^2} \left\{ v_{\parallel} (1 - c_s^2) \pm c_s \sqrt{(1 - v^2) \left[(1 - v^2 c_s^2) - v_{\parallel}^2 (1 - c_s^2) \right]} \right\}.\end{aligned}$$

The characteristic variables (corresponding to the left eigenvectors) are

$$\begin{aligned}l_{0,1} &= \frac{W}{1 - \mathcal{K}^{-1}} \{ h\sigma - W(\sigma + \rho_K) + W(v_{\parallel} J_{\parallel} + v_{\perp} J_{\perp} + \lambda^2 v^{\varphi} J^{\varphi}) \}, \\ l_{0,2} &= \frac{1}{h\xi} \left\{ -v_{\perp} (\sigma + \rho_K) + v_{\parallel} v_{\perp} J_{\parallel} + (1 - v_{\parallel}^2) J_{\perp} \right\}, \\ l_{0,3} &= \frac{1}{h\xi} \left\{ -v^{\varphi} (\sigma + \rho_K) + v^{\varphi} v_{\parallel} J_{\parallel} + (1 - v_{\parallel}^2) J^{\varphi} \right\}, \\ l_s^{\mp} &= \frac{h^2}{\Delta} \left\{ \mathcal{K}^{-1} hW \mathcal{V}^{\pm} \xi \sigma + [\mathcal{K}^{-1} - \mathcal{A}^{\pm} - (2 - \mathcal{K}^{-1}) v_{\parallel}] J_{\parallel} \right. \\ &\quad \left. + (2 - \mathcal{K}^{-1}) \mathcal{V}^{\pm} W^2 \xi (v_{\parallel} J_{\parallel} + v_{\perp} J_{\perp} + \lambda^2 v^{\varphi} J^{\varphi}) \right. \\ &\quad \left. + [(\mathcal{K}^{-1} - 1) (-v_{\parallel} + \mathcal{V}^{\pm} (W^2 \xi - 1)) - W^2 \mathcal{V}^{\pm} \xi] (\sigma + \rho_K) \right\}.\end{aligned}$$

The inverse transformation (corresponding to the right eigenvectors) is given by

$$\begin{aligned}\sigma &= \frac{1}{hW} l_{0,1} + W(v_{\perp} l_{0,2} + \lambda^2 v^{\varphi} l_{0,3}) + l_s^+ + l_s^-, \\ J_{\parallel} &= \mathcal{K}^{-1} v_{\parallel} l_{0,1} + 2hW^2 v_{\parallel} (v_{\perp} l_{0,2} + \lambda^2 v^{\varphi} l_{0,3}) + hW(\mathcal{C}^+ l_s^+ + \mathcal{C}^- l_s^-), \\ J_{\perp} &= \mathcal{K}^{-1} v_{\perp} l_{0,1} + h l_{0,2} + 2hW^2 v_{\perp} (v_{\perp} l_{0,2} + \lambda^2 v^{\varphi} l_{0,3}) + hW v_{\perp} (l_s^+ + l_s^-), \\ J^{\varphi} &= \mathcal{K}^{-1} v^{\varphi} l_{0,1} + h l_{0,3} + 2hW^2 v^{\varphi} (v_{\perp} l_{0,2} + \lambda^2 v^{\varphi} l_{0,3}) + hW v^{\varphi} (l_s^+ + l_s^-), \\ \rho_K &= \left(\mathcal{K}^{-1} - \frac{1}{hW} \right) l_{0,1} + W(2hW - 1)(v_{\perp} l_{0,2} + \lambda^2 v^{\varphi} l_{0,3}) \\ &\quad + hW(\mathcal{A}^+ l_s^+ + \mathcal{A}^- l_s^-) - l_s^+ - l_s^-.\end{aligned}$$

4. Transformation from conserved to primitive variables

The conserved matter variables (D1) are the ones that are evolved in a numerical algorithm. To compute the remaining matter variables (D7) and the eigenvectors, the primitive variables have to be calculated from the conserved variables as an intermediate step. This transformation is much more involved than the opposite direction (D6). To make it explicit, we have to specify an equation of state. Here, we consider the ultrarelativistic equation of state,

$$p = (\Gamma - 1) \rho_{tot} = (\Gamma - 1) \rho (\epsilon + 1) = \frac{\Gamma - 1}{\Gamma} \rho h, \quad (\text{D8})$$

where ρ_{tot} is the total energy density.

Suppose we are given the conserved variables, and also form $\rho_H = \rho_K + \sigma$. Consider the quantity

$$J^2 \equiv J_A J^A + \lambda^2 J^{\varphi 2}.$$

Using (D6), (D8) and (D5), we can express J^2 and ρ_H in terms of the primitive variables as

$$\begin{aligned}J^2 &= \left(\frac{\Gamma}{\Gamma - 1} \right)^2 p^2 W^2 (W^2 - 1), \\ \rho_H &= p \left(\frac{\Gamma}{\Gamma - 1} W^2 - 1 \right).\end{aligned}$$

Eliminating W yields an equation for the pressure in terms of conserved variables:

$$p = -2\beta \rho_H + \sqrt{4\beta^2 \rho_H^2 + (\Gamma - 1)(\rho_H^2 - J^2)},$$

where $\beta \equiv (2 - \Gamma)/4$. Next define

$$\chi_A \equiv \frac{(\Gamma - 1)J_A}{\Gamma p}, \quad \chi^\varphi \equiv \frac{(\Gamma - 1)J^\varphi}{\Gamma p}, \quad \chi^2 \equiv \chi_A \chi^A + \chi^\varphi \chi^\varphi.$$

We identify $\chi^A = W^2 v^A$ and $\chi^\varphi = W^2 v^\varphi$ and hence with (D5) we obtain

$$W^{-2} = \frac{1}{2\chi^2} \left(\sqrt{1 + 4\chi^2} - 1 \right). \quad (\text{D9})$$

This now enables us to calculate the velocities,

$$v_A = W^{-2} \chi_A, \quad v^\varphi = W^{-2} \chi^\varphi.$$

The form of W^{-2} in (D9) guarantees that $|v_A|, |v^\varphi| < 1$. This is most important since evolved speeds greater than unity (i.e. greater than the speed of light) can easily cause the numerical code to crash [16].

Finally, we can calculate the specific enthalpy and rest mass energy density from (D6) and (D8),

$$h = \frac{J^\varphi}{\sigma v^\varphi W}, \quad \rho = \frac{\Gamma p}{(\Gamma - 1)h}.$$

ACKNOWLEDGMENTS

We thank Anita Barnes for several useful discussions. OR thanks the Max Planck Institute for Gravitational Physics (Albert Einstein Institute), Golm, Germany for his inclusion in their Visitor Programme. OR acknowledges gratefully financial support from the Engineering and Physical Sciences Research Council, the Gates Cambridge Trust and Trinity College Cambridge.

-
- [1] Banyuls F, Font J A, Ibáñez J M, Martí J M and Miralles J A 1997 Numerical {3+1} General Relativistic Hydrodynamics: A Local Characteristic Approach *Astrophys. J.* **476** 221–231
 - [2] Bona C, Ledvinka T, Palenzuela C and Žáček M 2003 General-covariant evolution formalism for numerical relativity *Phys. Rev. D* **67** 104005
 - [3] Bona C, Ledvinka T, Palenzuela C and Žáček M 2004 Symmetry-breaking mechanism for the Z4 general-covariant evolution system *Phys. Rev. D* **69** 064036
 - [4] Bona C, Ledvinka T, Palenzuela-Luque C and Žáček M 2004 Constraint-preserving boundary conditions in the Z4 Numerical Relativity formalism *Preprint* arXiv:gr-qc/0411110
 - [5] Choptuik M W, Hirschmann E W, Liebling S L and Pretorius F 2003 An axisymmetric gravitational collapse code *Class. Quantum Grav.* **20** 1857–1878
 - [6] De Donder T 1921 *La Gravifique Einsteinienne* Gauthier-Villars, Paris
 - [7] Evans C R 1984 PhD. thesis, University of Texas, Austin, Texas, unpublished
 - [8] Fock V A 1959 *The Theory of Space Time and Gravitation* Pergamon, London
 - [9] Font J A 2003 Numerical Hydrodynamics in General Relativity *Living Rev. Rel.* **6** 4
 - [10] Garfinkle D and Duncan GC 2001 Numerical evolution of Brill waves *Phys. Rev. D* **63** 044011
 - [11] Geroch R 1971 A Method for Generating Solutions of Einstein's Equations *J. Math. Phys.* **12**(6) 918–924
 - [12] Gustafsson B, Kreiss H-O and Sundström A 1972 Stability Theory of Difference Approximations for Mixed Initial Boundary Value Problems. II *Math. Comput.* **26** 649–686
 - [13] Lehner L 2001 Numerical relativity: a review *Class. Quantum Grav.* **18** R25–R86
 - [14] Maeda K, Sasaki M, Nakamura T and Miyama S 1980 A New Formulation of the Einstein Equations for Relativistic Rotating Systems *Prog. Theor. Phys.* **63** 719–721
 - [15] Nakamura T, Oohara K and Kojima Y 1987 General Relativistic Collapse of Axially Symmetric Stars and 3D Time Evolution of Pure Gravitational Waves *Prog. Theor. Phys. Suppl.* **90** 13–109
 - [16] Nielsen D W and Choptuik M W 2000 Ultrarelativistic fluid dynamics *Class. Quantum Grav.* **17** 733–759
 - [17] Rauch J 1985 Symmetric positive systems with boundary characteristics of constant multiplicity *Trans. Am. Math. Soc.* **291** 167–187
 - [18] Secchi P 1996 The initial boundary value problem for linear symmetric hyperbolic systems with characteristic boundary of constant multiplicity *Diff. Int. Eq.* **9** 671–700, Well-posedness of characteristic symmetric hyperbolic systems *Arch. Rat. Mech. Anal.* **134** 155–197